First Hope

COMPLETE NOTES

Click here to visit our website

PHARMACY PRACTICAL NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Complete subjects and semesters are available in the FirstHope app.

Claisen-Schmidt Condensation

Practical 10 - Supervised synthesis and reaction-record practical

Aim

To prepare or assess dibenzalacetone by Claisen-Schmidt condensation and explain its reaction, isolation and characterisation.

Principle

Two benzaldehyde molecules and one acetone molecule form dibenzalacetone and two water molecules.

Base forms an acetone enolate; addition to the aldehyde gives a beta-hydroxy intermediate followed by dehydration. Repetition at the second methyl group extends conjugation.

Materials and Requirements

- Approved microscale reaction SOP and reagent specifications
- Supervised hood, temperature control, glassware and method-specific work-up equipment
- TLC references, balance and appropriate melting-point or spectral methods
- Approved method or teaching-data pack and a dated record sheet

Procedure

1. Define the preparation or dataset, study question and method version. Record identities, units and the criteria to be used before examining the outcome.
2. Draw the reaction and mechanism, identify stoichiometry, limiting reagent and possible side products. Enter the authorised quantities, addition order and temperature/time limits in the record.
3. Use the approved base-catalysed condensation procedure, controlling addition and mixing to limit competing reactions. Record precipitation and the TLC time course.
4. Monitor progress by the prescribed TLC or other analytical method, retaining the starting-material comparison.
5. Isolate and purify by the authorised work-up, record dry mass and examine identity and purity by complementary methods.
6. Retain the original observations and calculations. State deviations, missing information and the limits of the conclusion; do not replace measured values with expected results.

Observations and Record

Enter actual measurements or the supplied teaching data. Record the method, material identity, conditions and units.

Stage / preparation	Conditions / observation	Analytical evidence	Mass / yield
Starting materials	Identity / quantities	Reference evidence	Calculate mol
Reaction progress	Actual time / temperature	TLC or assay	Record
Purified product	Appearance	Identity / purity	Dry mass

Calculations

- Theoretical product mass = limiting-reactant moles x stoichiometric product ratio x product molar mass
- Isolated yield (%) = dry isolated product mass / theoretical product mass x 100
- Rf = zone migration / solvent-front migration

Worked Example

Illustrative 1:1 conversion from 10.0 mmol limiting reagent to a product of molar mass 234.29 g/mol gives 2.3429 g theoretical product. A dry isolated mass of 1.7572 g corresponds to 75.0% yield.

Use the actual reaction stoichiometry, salt or solvate form and purity basis; this illustration is not an expected experimental result.

Mechanism and identity record

• Base forms an acetone enolate; addition to the aldehyde gives a beta-hydroxy intermediate followed by dehydration. Repetition at the second methyl group extends conjugation.

• Compare the conjugated carbonyl and alkene evidence with the selected reference; note possible geometric isomers and residual benzaldehyde.

Result and Interpretation

• Report isolated yield and the independent identity/purity observations.

• Compare the conjugated carbonyl and alkene evidence with the selected reference; note possible geometric isomers and residual benzaldehyde.

Precautions

• Control caustic reagent and flammable solvent exposure.

• Use the institution-approved SOP and trained supervision for reactive additions, heating and work-up.

• Do not infer purity from yield or a single TLC zone.

Viva Questions and Answers

What determines theoretical yield?

The limiting reagent and balanced reaction stoichiometry.

How is conversion different from isolated yield?

Conversion measures reactant consumption; isolated yield also reflects selectivity and recovery losses.

Why use complementary identity tests?

A single physical or chromatographic observation may not distinguish impurities or isomers.

First Hope

COMPLETE NOTES

Click here to visit our website

PHARMACY PRACTICAL NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Complete subjects and semesters are available in the FirstHope app.