First Hope

COMPLETE NOTES

Click here to visit our website

PHARMACY PRACTICAL NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Complete subjects and semesters are available in the FirstHope app.

Preparation of Iodotoluene from Toluidine

Practical 15 - Supervised synthesis and reaction-record practical

Aim

To prepare or assess 4-iodotoluene from p-toluidine and explain its reaction, isolation and characterisation.

Principle

p-Toluidine is converted through an in-situ diazonium intermediate to 4-iodotoluene, retaining the para methyl relationship.

Diazotisation replaces the aromatic primary amine with a diazonium group; iodide substitution releases nitrogen. The methyl substitution position is retained.

Materials and Requirements

- Approved microscale reaction SOP and reagent specifications
- Supervised hood, temperature control, glassware and method-specific work-up equipment
- TLC references, balance and appropriate melting-point or spectral methods
- Approved method or teaching-data pack and a dated record sheet

Procedure

1. Define the preparation or dataset, study question and method version. Record identities, units and the criteria to be used before examining the outcome.
2. Draw the reaction and mechanism, identify stoichiometry, limiting reagent and possible side products. Enter the authorised quantities, addition order and temperature/time limits in the record.
3. Identify the toluidine isomer and examine the authorised cold, in-situ diazotisation and immediate substitution record. Use only the supervised approved scale or a supplied dataset; record the defined temperature limits and work-up.
4. Monitor progress by the prescribed TLC or other analytical method, retaining the starting-material comparison.
5. Isolate and purify by the authorised work-up, record dry mass and examine identity and purity by complementary methods.
6. Retain the original observations and calculations. State deviations, missing information and the limits of the conclusion; do not replace measured values with expected results.

Observations and Record

Enter actual measurements or the supplied teaching data. Record the method, material identity, conditions and units.

Stage / preparation	Conditions / observation	Analytical evidence	Mass / yield
Starting materials	Identity / quantities	Reference evidence	Calculate mol
Reaction progress	Actual time / temperature	TLC or assay	Record
Purified product	Appearance	Identity / purity	Dry mass

Calculations

- Theoretical product mass = limiting-reactant moles x stoichiometric product ratio x product molar mass
- Isolated yield (%) = dry isolated product mass / theoretical product mass x 100
- Rf = zone migration / solvent-front migration

Worked Example

Illustrative 1:1 conversion from 10.0 mmol limiting reagent to a product of molar mass 218.04 g/mol gives 2.1804 g theoretical product. A dry isolated mass of 1.6353 g corresponds to 75.0% yield.

Use the actual reaction stoichiometry, salt or solvate form and purity basis; this illustration is not an expected experimental result.

Mechanism and identity record

- Diazotisation replaces the aromatic primary amine with a diazonium group; iodide substitution releases nitrogen. The methyl substitution position is retained.
- Verify isomer identity by suitable spectra and chromatographic comparison rather than assuming all iodotoluenes are equivalent.

Result and Interpretation

- Report isolated yield and the independent identity/purity observations.
- Verify isomer identity by suitable spectra and chromatographic comparison rather than assuming all iodotoluenes are equivalent.

Precautions

- Do not isolate, dry, store or concentrate diazonium salts. The supervised in-situ process requires explicit thermal and gas controls.
- Use the institution-approved SOP and trained supervision for reactive additions, heating and work-up.
- Do not infer purity from yield or a single TLC zone.

Viva Questions and Answers

What determines theoretical yield?

The limiting reagent and balanced reaction stoichiometry.

How is conversion different from isolated yield?

Conversion measures reactant consumption; isolated yield also reflects selectivity and recovery losses.

Why use complementary identity tests?

A single physical or chromatographic observation may not distinguish impurities or isomers.

First Hope

COMPLETE NOTES

Click here to visit our website

PHARMACY PRACTICAL NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Complete subjects and semesters are available in the FirstHope app.