First Hope

COMPLETE NOTES

Click here to visit our website

PHARMACY PRACTICAL NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Complete subjects and semesters are available in the FirstHope app.

Raw-Material Analysis Documentation

Practical 3 - Analytical practical

Aim

To plan and document the official assay and identification evidence required for a raw material.

Principle

A titration relates analyte amount to the amount of standard reagent required by a defined reaction. Reaction stoichiometry, endpoint selection, blank consumption and reagent standardisation determine the calculation. The test preparation, blank, calibration or titrant standardisation and independent controls are part of the measurement. State the actual concentration units, test conditions and adopted method before interpreting the result.

Materials and Requirements

- Selected official compound and adopted monograph
- Appropriate standardised titrant and primary-standard record
- Endpoint indicator or potentiometric system
- Qualified balance and volumetric equipment
- Controlled method and raw-data worksheet

Procedure

1. Identify the material, intended attribute, method version and acceptance criteria. Record preparation and instrument or reagent identities.
2. Prepare the prescribed blank and reference or standardised reagent. Verify the measurement with an independent check where applicable.
3. Prepare a representative weighed test portion in the specified solvent. Ensure the reaction conditions and dissolution are appropriate.
4. Titrate using the defined endpoint, recording initial and final burette readings and a reagent blank.
5. Repeat on independent portions and calculate the content using the reaction stoichiometry and the actual titrant concentration.
6. Record individual readings for independently prepared test portions. Keep all raw observations, including unsuccessful measurements and their investigation.
7. Calculate the result on the stated mass, volume or area basis, including dilution, blank and recovery corrections as justified.
8. Compare the evidence with the applicable criteria and write a conclusion limited to the tested preparation and conditions.

Observations and Record

Enter actual measurements or the supplied teaching data. Record the method, material identity, conditions and units.

Preparation / attribute	Method condition	Raw measurement	Calculated result
Titrant standardisation	Record	Record	Calculate
Blank	Record	Record	Calculate
Test portion 1	Record	Record	Calculate

Preparation / attribute	Method condition	Raw measurement	Calculated result
Test portion 2	Record	Record	Calculate

Calculations

• Analyte amount (mol) = blank-corrected titrant volume (L) x actual titrant molarity x stoichiometric ratio
• Assay (%) = measured analyte mass / weighed preparation mass x 100

Worked Example

Illustrative 1:1 reaction: corrected titrant volume 10.00 mL, molarity 0.1000 mol/L and analyte molar mass 180.16 g/mol.

Measured analyte = 0.01000 x 0.1000 x 180.16 = 0.18016 g.

For a 0.1820 g test portion, assay = 98.99% approximately. This example does not assign the reaction to a particular official drug.

Monograph-to-record mapping

• Identify the current applicable monograph, material grade and any water, solvent or purity correction.
• Map identity, assay and impurity requirements to method records and raw data. A supplier certificate supports the record but does not replace required receipt testing.

Result and Interpretation

• State the endpoint and stoichiometric basis, including whether water or other correction factors are required by the monograph.
• Report replicate variability and the evidence that the method can measure the intended attribute in this matrix.

Precautions

• Use the actual standardised reagent concentration.
• Avoid endpoint overshoot and parallax.
• Select a blank with the same reagent and solvent conditions.
• Use the adopted laboratory procedure, suitable controls and the designated waste arrangements.

Viva Questions and Answers

Why standardise a titrant?

Its actual concentration can differ from the nominal preparation.

Why is stoichiometry essential?

The reagent-to-analyte mole relationship determines the mass calculation.

Why correct for the blank?

Other components may consume titrant.

First Hope

COMPLETE NOTES

Click here to visit our website

PHARMACY PRACTICAL NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Complete subjects and semesters are available in the FirstHope app.