First Hope

COMPLETE NOTES

Click here to visit our website

PHARMACY PRACTICAL NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Complete subjects and semesters are available in the FirstHope app.

Preparation and Evaluation of Alginate Beads

Practical 2 - Bench formulation practical

Aim

To prepare calcium-alginate beads and evaluate formation, size, recovery and marker release.

Principle

Alginate chains form an ionically cross-linked gel in the presence of suitable calcium ions. Droplet formation controls initial size, while polymer grade, solution viscosity, cross-linking and subsequent handling affect the bead structure.

A formulation comparison is meaningful when the raw-material grades, batch size and evaluation conditions are recorded. Change one planned factor at a time or use a documented experimental design, and compare independently prepared batches rather than only repeated readings from one batch.

Materials and Requirements

- Sodium alginate and calcium chloride solutions prepared by the approved method
- Benign marker or approved model drug
- Controlled dispensing device, stirrer and bead-sizing method
- Calibrated balance, suitable containers and identified evaluation equipment
- Controlled batch sheet and raw-data forms

Procedure

1. Define the purpose, formula, batch size, material grades and comparison conditions. Calculate all quantities before weighing.
2. Check equipment readiness and record material identities and actual quantities. Reconcile theoretical and actual batch mass.
3. Hydrate the selected alginate grade and document its concentration, viscosity condition and marker content.
4. Introduce controlled droplets into the prescribed calcium bath with the approved nozzle and mixing conditions.
5. Maintain the chosen curing procedure, then recover and rinse the beads using the stated method.
6. Measure wet size and the defined dry or wet recovery basis. Assess marker content and release using validated separation and assay steps.
7. Evaluate the stated quality attributes on representative portions using the same method for all comparison batches. Record individual readings and any excluded observations with reasons.
8. Calculate the specified response, variability and material balance. Compare with predefined teaching or product criteria and discuss sources of variation.
9. Label retained preparations and dispose of remaining material through the laboratory procedure. Keep the complete preparation and evaluation record together.

Observations and Record

Enter actual measurements or the supplied teaching data. Record the method, material identity, conditions and units.

Attribute or stage	Method / conditions	Individual readings	Calculation / comment
Droplet and bath conditions	Record the adopted method	Record	Evaluate

Attribute or stage	Method / conditions	Individual readings	Calculation / comment
Bead size distribution	Record the adopted method	Record	Evaluate
Recovered mass	Record the adopted method	Record	Evaluate
Marker content and release	Record the adopted method	Record	Evaluate

Calculations

• Encapsulation efficiency (%) = entrapped marker / input marker x 100
• Drug loading (%) = entrapped marker / recovered dry bead mass x 100

Worked Example

Illustrative input marker is 100 mg and the validated measurement of free marker is 20 mg.
Entrapped marker is 80 mg and encapsulation efficiency is 80.0%. If recovered dry beads weigh 400 mg, loading is 20.0% w/w.
Account for recovery and assay losses before assigning all unmeasured marker to encapsulation.

Result and Interpretation

• Relate bead size and release to the actual process conditions. State whether mass and size were measured wet or dry.
• State the preparation conditions and the evidence supporting the result. A satisfactory physical observation alone does not establish long-term stability, clinical performance or suitability for administration.

Precautions

• Control the dispensing geometry and solution condition.
• Prevent contamination during curing and recovery.
• Do not compare wet-yield and dry-yield percentages as equivalent quantities.
• Use the laboratory-approved formulation and equipment procedure. Teaching batches are identified for laboratory evaluation.

Viva Questions and Answers

What forms the gel?

Ionic cross-linking of alginate by calcium ions.

Why record alginate grade?

Polymer characteristics influence viscosity and gel behaviour.

Are loading and encapsulation efficiency identical?

No; they use different denominator quantities.

First Hope

COMPLETE NOTES

Click here to visit our website

PHARMACY PRACTICAL NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Complete subjects and semesters are available in the FirstHope app.