

Benzoylation - Benzanilide and Phenyl Benzoate

Experiment 2 - Practical Notes

Name: _____ Roll no.: _____ Date: _____

Aim

To study the specified organic preparation, isolate the assigned product and calculate its yield.

Principle

Benzoylation introduces a benzoyl group. Amines form amide derivatives and phenols can form ester derivatives.

Requirements

- Instructor-approved reaction scheme, reagent quantities and laboratory procedure for the selected route.
- Reaction glassware, suitable ventilation, separation apparatus, balance and melting-point apparatus.

Procedure

1. Plan the selected route

Write the balanced reaction and identify the functional-group change, limiting reagent, expected product and hazards. Use the assigned supervised method for the selected option.

2. Carry out the assigned reaction

Measure the prescribed reagents and follow the authorised addition sequence and temperature controls. Record actual reaction observations and the method used.

3. Isolate the product

Separate the product using the assigned acid/base work-up, filter the solid and wash until the specified residual reagents are removed.

4. Purify and dry

Apply the assigned recrystallisation or purification method. Record solvent, observations, dry mass and melting range where appropriate.

5. Assess the product

Calculate theoretical and percentage yield. Compare the measured physical properties and assigned identity test; explain that yield alone does not prove purity.

Preparations and Reaction Scheme

Route / product	Starting material and change
Benzanilide	Aniline to the corresponding benzamide derivative.
Phenyl benzoate	Phenol to a benzoate ester.

Pharmaceutical Organic Chemistry

Observations and Record

Record	Student entry
Selected route / balanced equation	_____
Limiting reagent and molar mass	_____
Theoretical product mass	_____
Dry product mass / melting range	_____
Percentage yield / identity finding	_____

Worked Example

The numbers below are illustrative; enter your own observations in the record.

- Theoretical product mass = moles of limiting reagent $\times$ stoichiometric product ratio $\times$ product molar mass.
- Percentage yield = dry product mass / theoretical product mass $\times$ 100.
- Illustration: 0.020 mol of a limiting reagent gives 0.020 mol of a 150 g/mol product. The theoretical mass is 3.00 g; 2.40 g isolated product gives 80.0% yield.

Result

Product: _____. Route: _____. Dry mass: _____. Yield: _____. Physical property / identity evidence: _____.

Precautions

- Use the specific supervised procedure and facilities for corrosive, toxic, oxidising or flammable reagents.
- Record the selected preparation option rather than combining different reaction routes.
- Use dry product mass for yield and the stated test conditions for comparison.

Viva Questions and Answers**1. Why does reaction-site identity matter?**

Nitrogen and oxygen give different derivative classes.

2. What is the purpose of work-up?

It separates the desired product from residual reagents and by-products.

Continue Learning with FirstHope

Complete Notes on the FirstHope App

Complete subject and semesters are available on FirstHope app only.

- Get complete subject and semester notes in the FirstHope app.
- Available for B.Pharm, D.Pharm, M.Pharm and Pharm.D students.

[Click here to get the FirstHope app](#)

Join the FirstHope Telegram Channel

Join FirstHope Notes and keep up with channel updates.

[Click here to join Telegram](#)

Instagram

@firsthope.official

[Click here to follow Instagram](#)

YouTube Channel

FirstHope-Pharmacy / Live streams

[Click here to watch the YouTube channel](#)

WhatsApp Channel

Follow FirstHope on WhatsApp.

[Click here to join the WhatsApp channel](#)

FirstHope Website

firsthopenotes.com

[Click here to visit the website](#)