

Dietary Supplements and Nutraceuticals

Effect of Processing on the Potential of Nutraceuticals

- Processing plays a vital role in determining the bioavailability, efficacy, and stability of nutraceuticals.
- While some processing techniques can enhance these properties, others may degrade the active compounds if not carefully managed.

Impact on Bioavailability

A. Extraction Methods

- Techniques like cold pressing and solvent extraction can improve the availability of bioactive compounds by making them more accessible for absorption.
- However, harsh methods may degrade heat-sensitive or light-sensitive components.

B. Encapsulation

- Encapsulation protects active ingredients from degradation during storage and digestion, enhancing both bioavailability and targeted delivery.

Influence on Efficacy

A. Thermal Processing

- High heat can denature or destroy delicate bioactive compounds, reducing therapeutic potential.
- In contrast, heat may sometimes activate or convert compounds into more bioactive forms (e.g., lycopene in tomatoes).

B. Fermentation

- Fermentation may improve the efficacy of nutraceuticals by enhancing digestibility and producing bioactive metabolites with improved biological activity.

Effects on Stability

A. Drying Techniques

- Methods like freeze-drying and spray drying help reduce moisture, improving shelf life and preserving compound integrity.

B. Microencapsulation

- This process offers a protective barrier against oxidative and environmental damage, enhancing product stability during storage.

Modification of Active Components

A. Chemical Modifications

- Some processing methods may chemically alter active ingredients, creating new compounds with different or reduced biological activity.

B. Degradation of Nutrients

- Excessive processing can lead to the breakdown of essential nutrients, reducing the nutritional value and effectiveness of the product.

Effect of Storage on the Potential of Nutraceuticals

- Storage conditions greatly influence the safety, stability, and potency of nutraceuticals.
- Factors like temperature, light, humidity, oxygen exposure, and packaging play key roles in preserving product quality.

1. Temperature

A. Heat Exposure

- High temperatures accelerate degradation of sensitive compounds such as antioxidants and vitamins, leading to reduced efficacy.

B. Cold Storage

- Refrigeration can slow degradation and extend shelf life.
- However, extreme cold may cause precipitation or destabilization of some ingredients.

2. Light Exposure

A. UV and Visible Light

- Light exposure can lead to photooxidation and degradation of compounds like riboflavin, vitamin A, and phytochemicals.

B. Protective Packaging

- Using opaque, UV-resistant containers helps prevent light-induced deterioration.

3. Humidity

A. Moisture Absorption

- High humidity can lead to clumping, microbial contamination, and hydrolysis, all of which affect product integrity.

B. Use of Desiccants

- Incorporating desiccants in packaging can help control internal moisture levels and protect hygroscopic compounds.

4. Oxygen Exposure

A. Oxidation

- Oxygen can degrade sensitive nutrients such as vitamin C, vitamin E, and polyunsaturated fatty acids, reducing their effectiveness and possibly forming harmful by-products.

B. Oxygen Scavengers & Inert Packaging

- Techniques like nitrogen flushing and oxygen scavengers reduce oxidation by minimizing oxygen exposure.

5. Interaction with Packaging Materials

A. Material Reactivity

- Packaging materials can interact chemically with the product, leading to contamination or degradation.
- Selecting inert, non-reactive packaging is crucial to avoid such interactions.

Effect of interactions of various environmental factors on the potential of nutraceuticals

- The potential of nutraceuticals—defined by their bioavailability, stability, and efficacy—can be significantly influenced by environmental interactions.
- These complex interactions involve variables such as temperature, humidity, light, oxygen, pollutants, and even packaging materials.
- Understanding these interdependent effects is crucial to preserving the quality and therapeutic potential of nutraceuticals.

1. Temperature and Humidity Interactions

A. Moisture-Induced Degradation

- Elevated temperatures can enhance moisture absorption in hygroscopic materials, accelerating degradation reactions like hydrolysis.

B. Thermal Stability in Humid Conditions

- Some compounds remain stable at high temperatures under dry conditions but degrade rapidly when exposed to both heat and moisture simultaneously.

2. Light and Oxygen Exposure

A. Photooxidation

- UV and visible light in the presence of oxygen can trigger oxidative reactions, degrading sensitive compounds such as:
 - Vitamins (e.g., riboflavin, vitamin A)
 - Antioxidants
 - Unsaturated fatty acids




FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.