Hyphenated techniques

Introduction to Hyphenated Techniques

- Hyphenated techniques are analytical methods that combine two or more instrumental techniques to leverage the strengths of each.
- Typically, this involves chromatographic separation (like LC, GC, or HPTLC) followed by spectroscopic detection (like MS, UV, or IR).
- These techniques provide:
    - Separation (via chromatography)
    - Identification (via mass spectrometry or spectroscopy)
    - Quantification (based on peak area or intensity)

Purpose:

- Improve sensitivity, selectivity, structural elucidation, and quantification in complex sample matrices.

Examples of Hyphenated Techniques:

- **LC-MS/MS** – Liquid Chromatography–Tandem Mass Spectrometry
- **GC-MS/MS** – Gas Chromatography–Tandem Mass Spectrometry
- **HPTLC-MS** – High Performance Thin Layer Chromatography–Mass Spectrometry

LC-MS/MS (Liquid Chromatography – Tandem Mass Spectrometry)

General Principle:

- LC separates analytes in a liquid mobile phase based on polarity, size, or charge.
- MS/MS (Tandem Mass Spectrometry) involves two stages of mass analysis:
    - The first MS isolates the parent ion (precursor).
    - The second MS analyzes the fragmented ions (daughter ions).
- Electrospray Ionization (ESI) or APCI is typically used to interface LC with MS.

Components

Component	Function
Liquid Chromatograph (LC)	Separates components in a liquid phase using a column and mobile phase.

Ionization Source	Converts eluted molecules into gas-phase ions (commonly ESI – Electrospray Ionization, or APCI – Atmospheric Pressure Chemical Ionization).
Mass Spectrometer (Triple Quadrupole)	Detects ions and fragments them to produce structural data. Consists of Q1 – Q2 – Q3.

Procedure / Workflow:

1. **Sample Injection into LC**
   - Components are separated based on interaction with column.
2. **Ionization (ESI/APCI)**
   - Converts eluted analytes into gas-phase ions.
3. **MS1 (Precursor Ion Selection)**
   - Selects a specific ion of interest.
4. **Collision Cell (CID)**
   - Precursor ion is fragmented using inert gas (e.g., nitrogen).
5. **MS2 (Product Ion Analysis)**
   - Detects specific fragment ions.
6. **Data Interpretation**
   - Quantification and identification using MRM (Multiple Reaction Monitoring).

Advantages:

- Extremely high sensitivity and specificity
- Can detect picogram to femtogram levels
- Excellent for complex biological matrices (plasma, serum, urine)
- Multiple analytes can be quantified simultaneously

Applications:

- **Pharmaceuticals**: Drug metabolism studies, bioequivalence.
- **Clinical**: Hormone quantification (e.g., testosterone, vitamin D).
- **Toxicology**: Drugs of abuse, forensic investigations.
- **Food safety**: Pesticide residue analysis.
- **Environmental**: Pollutants in water and soil.

GC-MS/MS (Gas Chromatography – Tandem Mass Spectrometry)

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.