

Clinical Trials

Clinical Trials: Types and Design

- Clinical trials are research studies done in humans to find out whether a drug, vaccine, device, treatment, or medical procedure is safe and effective.



- Clinical trials help researchers understand how well a treatment works, what side effects it may cause, and whether it is better than the current treatment.

Basic Design of a Clinical Trial

A. Research Question

- The study begins with a clear question.
- Example: Researchers may ask whether a new drug lowers blood pressure better than an existing drug.

B. Study Population

- The study population means the group of people selected for the trial.
- These people may have a particular disease or health condition.

C. Inclusion Criteria

- Inclusion criteria are the conditions a person must meet to enter the study.
- Example: A study may include adults between 18 and 60 years of age with diabetes.

D. Exclusion Criteria

- Exclusion criteria are the conditions that prevent a person from joining the study.
- Example: A study may exclude pregnant women or patients with severe kidney disease.

E. Intervention

- An intervention is the treatment given during the trial.
- The intervention may be a drug, vaccine, surgery, diet plan, or medical device.

F. Control Group

- A control group is used for comparison.
- The control group may receive a placebo, standard treatment, or no treatment.

G. Randomization

- Randomization means participants are placed into groups by chance.
- This helps reduce bias and makes the groups more equal.

H. Blinding

- Blinding means hiding treatment information from participants, researchers, or both.
- Blinding helps prevent personal opinions from affecting the results.


I. Outcome Measurement

- Outcomes are the results measured in the study.
- Example: Blood pressure, survival rate, pain relief, or side effects can be outcomes.

J. Follow-Up

- Follow-up means checking participants after treatment for a certain period.
- Follow-up helps researchers see both short-term and long-term effects.

Types of Clinical Trials



1. Experimental Clinical Trials

- In experimental clinical trials, researchers give an intervention and study its effect.
- These trials include randomized controlled trials and non-randomized trials.

2. Observational Clinical Studies

- In observational studies, researchers do not give any treatment.
- They only observe and collect information from participants.

3. Randomized Trials

- In randomized trials, participants are assigned to groups by chance.
- This type of trial gives strong evidence about treatment effects.

4. Non-Randomized Trials

- In non-randomized trials, participants are not assigned by chance.
- The researcher or situation decides which group the participant enters.

5. Blinded Trials

- In blinded trials, participants or researchers do not know which treatment is being given.
- This helps reduce bias in the study.

6. Open-Label Trials

- In open-label trials, both participants and researchers know which treatment is being given.
- These trials are easier to conduct but may have more bias.

Experimental Study

- An experimental study is a study in which the researcher actively gives an intervention and observes its effect.
- The intervention may be a medicine, vaccine, surgery, health education program, or any treatment method.

Main Features of Experimental Study

- The researcher controls the intervention.
- The participants are divided into groups.
- One group usually receives the treatment.
- Another group may receive standard treatment, placebo, or no treatment.
- The results of both groups are compared.
- Experimental studies are useful for finding cause-and-effect relationships.

Randomized Controlled Trial

- A randomized controlled trial is an experimental study in which participants are divided into groups by chance.
- Randomized controlled trials are considered strong studies because randomization reduces bias.


Important Features of RCT

- Participants are randomly assigned to different groups.
- One group receives the new treatment.
- Another group receives standard treatment, placebo, or no treatment.
- The outcomes of both groups are compared.
- Randomization helps make the groups similar before treatment begins.

Example of RCT

- A group of patients with high blood pressure is selected.
- Half of the patients receive a new blood pressure medicine.
- The other half receive the standard medicine.
- After some time, researchers compare blood pressure changes in both groups.

Subtypes of RCT



A. Parallel RCT

- In a parallel RCT, each participant stays in one treatment group throughout the study.
- One group receives treatment A, and another group receives treatment B.

B. Crossover RCT

- In a crossover RCT, each participant receives more than one treatment at different times.
- The same person may receive treatment A first and treatment B later.

C. Cluster RCT

- In a cluster RCT, groups of people are randomized instead of individual persons.
- Example: Entire schools, villages, or hospitals may be assigned to different interventions.

D. Factorial RCT

- In a factorial RCT, researchers study two or more interventions at the same time.
- This design helps check the effects of separate treatments and combined treatments.

Advantages of Randomized Controlled Trials (RCTs)

- RCTs provide strong and reliable evidence about the effectiveness of a treatment or intervention.
- Random assignment of participants helps minimize selection bias and ensures fair group allocation.
- RCTs enable clear and direct comparisons between treatment and control groups.

Limitations of Randomized Controlled Trials (RCTs)

- Conducting RCTs can be costly and time-consuming.
- RCTs may be unethical or impractical when studying potentially harmful exposures.
- Some individuals may decline participation because they cannot choose which treatment they receive.

Non-Randomized Controlled Trial

- A non-randomized controlled trial is an experimental study in which participants are not assigned to groups by chance.
- In this type of study, the researcher still gives an intervention, but the group assignment is decided by other methods.

Important Features of Non-RCT

- Participants are not randomly assigned.
- The researcher may decide which group receives the treatment.
- The study may compare treated patients with untreated patients.
- This study is easier to conduct than an RCT.
- This study has a higher chance of bias than an RCT.

Example of Non-RCT

- One hospital starts using a new treatment for asthma.
- Another hospital continues using the old treatment.
- Researchers compare patient improvement in both hospitals.



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.