Natural Products as Leads for New Pharmaceuticals

Introduction to Natural-Product Drug Discovery

- A natural-product lead supplies a biologically active chemical framework that can be studied directly or modified to improve useful properties.
    - A semisynthetic derivative retains part of a naturally obtained molecule and changes selected groups through chemical reactions.

What a Natural Lead Can Contribute

- A **lead** is a compound with useful, reproducible biological activity that provides a starting point for further chemical and pharmaceutical improvement.
- Natural products often contain defined three-dimensional frameworks and several functional groups that recognise biological macromolecules.
    - A natural compound may require modification to improve supply, solubility, selectivity, stability or delivery.

Semisynthesis and Total Synthesis

- **Semisynthesis** modifies a naturally obtained starting material, whereas total synthesis constructs the target from simpler chemical precursors.
- A natural origin does not establish safety, clinical efficacy or suitability for every formulation.
    - The study of a natural lead should connect its source, chemical class, important structural features and the reason that analogues were developed.
- A successful lead programme can preserve a complex natural scaffold, simplify it or translate only its key interaction pattern into another chemical class.
    - The optimal strategy depends on whether complexity is essential for binding and whether the material can be supplied consistently.
- Natural-product identification, semisynthesis, target studies and process development therefore contribute different parts of the discovery problem.

Natural Products as Leads for New Pharmaceuticals

Three Development Routes

- The natural compound itself may become a medicine when its properties and supply are suitable.
- A semisynthetic derivative retains much of the natural framework but changes selected groups.
- A simpler synthetic analogue may reproduce only the essential features of the lead.

First Hope

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

Evidence at Each Stage

- Identity, purity and reproducible activity must be established before interpreting a natural extract.
    - An effect of a mixture cannot automatically be assigned to its most abundant constituent.

Examples of Natural-Lead Development

Natural lead or family	Main development problem	Medicinal-chemistry response
Teprotide and related peptides	Peptide-like properties limited oral usefulness.	Smaller ACE-inhibitor scaffolds retained selected recognition features.
Erythromycin A	Acid instability and disposition limited some uses.	Selective methylation, oxime modification or ring expansion gave related macrolides.
Taxanes	Complexity, supply and poor aqueous solubility complicated development.	Semisynthetic precursors, side-chain modification and formulation work supported use.
Lovastatin	The initial natural lead provided room for potency and disposition changes.	Modification of the hydrophobic region produced related statin designs.
Podophyllotoxin-related lignans	The original scaffold did not provide the desired anticancer profile.	Semisynthetic changes produced etoposide and teniposide with a different primary target mechanism.

A Natural-Lead Evaluation Framework

- Lead evaluation connects a defined chemical structure with activity, exposure and limitations. A promising assay result is followed by comparison of modified compounds using the same relevant measurements.



Evaluation Sequence

A. Establish Activity and the Limiting Property

- First establish that the biological activity belongs to the identified compound and is reproducible.

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

- Next determine the important structural features and identify a limiting property such as supply, selectivity or stability.

B. Modify and Compare

- Choose a modification that addresses that limitation while preserving the productive binding arrangement.
  - Finally compare the analogue with the parent using matched biological and physicochemical measurements.

Morphine Alkaloids as Central Nervous System Leads

Systematic Study

- **Morphine** is an alkaloid associated with opium poppy and contains a rigid, bridged polycyclic framework with a basic nitrogen.


Structure–Activity Relationships

Key Structural Features

A. Aromatic and Oxygen-Containing Groups

- Morphine contains an aromatic ring, a phenolic hydroxyl group, an alcoholic hydroxyl group and an ether bridge in a rigid polycyclic framework.
  - The phenolic group contributes to receptor recognition, while changing an oxygen-containing group can alter lipophilicity, metabolism and activity.
- Methylation of the phenolic oxygen gives codeine.
  - Codeine's analgesic effect partly depends on metabolic conversion to morphine, so variation in metabolism has clinical importance.

First Hope



Modification of an oxygen substituent changes the properties of a closely related alkaloid scaffold

B. Basic Nitrogen and Stereochemistry

- The tertiary amine is substantially protonated under physiological conditions and can participate in ionic recognition.

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

        - Changing the nitrogen substituent can change receptor efficacy as well as affinity.
    - The three-dimensional arrangement of the rings and substituents is essential; a formula alone does not define the active pharmacophore.
        - An increase in lipophilicity can change access to the central nervous system without automatically improving the overall benefit-risk profile.

Mechanism of Action

- Morphine is an opioid analgesic whose major actions arise from activation of μ-opioid receptors.
- These receptors couple mainly to inhibitory G proteins, reducing neuronal transmitter release and excitability.
- Analgesia and adverse effects arise from receptor actions in different neural circuits; a stronger analgesic effect does not remove the risks of respiratory depression and dependence.

What the Lead Teaches

- Changes around the nitrogen or oxygen-containing groups can alter receptor affinity, intrinsic activity and disposition.
    - A structurally related opioid can be an agonist, partial agonist or antagonist depending on the complete recognition pattern.
- The class also illustrates serious limitations related to dependence, respiratory depression and other unwanted effects.
    - Structural resemblance should therefore not be equated with identical pharmacological behaviour.

First Hope

Value and Limitations of the Lead

- Morphine-related chemistry illustrates the separation of analgesic activity from undesirable effects such as respiratory depression, constipation and dependence.
- A structural modification must be assessed experimentally because improved potency does not automatically mean an improved therapeutic margin.


Paclitaxel as an Anticancer Lead

Recognising the Taxane Framework

- Paclitaxel has a complex taxane core, an oxetane ring and an important side chain attached through an ester linkage.
    - Its stereochemistry helps organise the groups required for microtubule binding.
- **Paclitaxel** contains more than twenty carbon atoms because its diterpenoid core carries additional substituents.

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

Paclitaxel



Paclitaxel

Paclitaxel and docetaxel share a taxane framework but differ in substituents — Paclitaxel

Mechanism of Action

- **Paclitaxel** binds to microtubules and favours their stable assembled state. Suppression of normal microtubule dynamics disrupts mitotic spindle function and cell division.

Lead-Optimisation Lessons

- The natural scaffold supplies strong biological recognition but creates challenges in solubility, formulation and supply.
    - Semisynthetic approaches can use an appropriate naturally derived core and attach a suitable side chain.
- A modification must preserve the essential binding arrangement while improving a useful pharmaceutical property.

Semisynthetic Supply

- **Docetaxel** is a related semisynthetic taxane developed through modification of an accessible taxane precursor.
    - The naturally obtainable intermediate 10-deacetylbaccatin III supports semisynthetic supply strategies for taxane medicines.
    - Taxane development illustrates the combined importance of a biological lead, renewable precursor supply, side-chain chemistry and formulation.

Docetaxel as a Semisynthetic Taxane

Structural Differences and Medicinal Consequences

A. Shared Taxane Core

- Both agents retain a taxane framework and an ester-linked side chain at C13.

First Hope

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.