Preclinical screening of Respiratory Pharmacology

Anti-Asthmatics – Preclinical Screening Methods

- **Asthma is a chronic inflammatory disease of airways characterized by:**
  - Bronchoconstriction (narrowing of airways)
  - Inflammation
  - Mucus secretion
- **Anti-asthmatic drugs work by:**
  - Dilating bronchi (bronchodilators)
  - Reducing inflammation (anti-inflammatory drugs)
  - Preventing allergic responses
- Preclinical screening identifies substances that reduce airway resistance and inflammation.

Objectives of Screening

- To evaluate bronchodilatory activity
- To assess anti-inflammatory effects in airways
- To study mechanism (β2 agonist, leukotriene inhibition, etc.)
- To detect toxicity (especially cardiac and respiratory effects)

First Hope

In Vivo Methods (Animal Models)

A. Histamine-Induced Bronchospasm Model

- **Principle:** Histamine causes bronchoconstriction.
- **Method:**
  - Expose guinea pigs to histamine aerosol
  - Measure time to onset of dyspnea (difficulty breathing)
- **Observation:** Increased time before dyspnea = bronchodilator effect
- **Importance:** Mimics acute asthma attack

B. Acetylcholine-Induced Bronchospasm

- **Principle:** Acetylcholine causes airway constriction via muscarinic receptors
- **Method:** Aerosol administration of acetylcholine
- **Observation:** Protection against bronchospasm = anticholinergic or bronchodilator activity

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

C. Ovalbumin-Induced Allergic Asthma Model

- **Principle:** Mimics allergic asthma using antigen exposure
- **Method:**
  - Sensitize animals (usually mice) with ovalbumin
  - Re-expose to trigger airway inflammation
- **Observation:** Reduced airway inflammation, eosinophils, mucus = anti-asthmatic effect

D. Measurement of Airway Resistance

- **Method:** Use plethysmography to measure lung function
- **Observation:** Decreased airway resistance = bronchodilation

E. Passive Cutaneous Anaphylaxis (PCA)

- **Principle:** Measures of allergic response mediated by IgE
- **Observation:** Reduced vascular permeability = anti-allergic/anti-asthmatic effect

In Vitro Methods

1. Isolated Tracheal Chain Preparation

- **Method:** Tracheal rings from animals mounted in organ bath
- **Observation:** Relaxation of pre-contracted trachea = bronchodilator activity

2. Mast Cell Stabilization Assay

- **Principle:** Mast cells release histamine during asthma
- **Method:** Induce degranulation in mast cells
- **Observation:** Inhibition of histamine release = anti-asthmatic effect

3. Leukotriene Assays

- Measure inhibition of leukotriene production (important inflammatory mediators)

Other Alternative Models

- **Zebrafish models:** Used for inflammation and respiratory movement
- **Cell cultures:** Airway epithelial cells to study cytokine release

Parameters Measured

- Airway resistance
- Bronchospasm severity
- Inflammatory cell count (eosinophils)

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.


- Cytokine levels

Toxicological Considerations

- Cardiac stimulation (β2 agonists)
- Tremors
- Long-term steroid effects (if anti-inflammatory drugs)

Drugs for COPD (Chronic Obstructive Pulmonary Disease) – Preclinical Screening Methods

- **COPD is a progressive lung disease characterized by:**
  - Chronic bronchitis (mucus hypersecretion)
  - Emphysema (destruction of alveoli)
  - Persistent airflow limitation



- **Drugs aim to:**
  - Dilate airways
  - Reduce inflammation
  - Improve mucus clearance

Objectives of Screening

- To evaluate bronchodilation
- To assess anti-inflammatory effects
- To study mucus secretion control
- To model chronic lung damage

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

In Vivo Methods (Animal Models)

A. Cigarette Smoke-Induced COPD Model

- **Principle:** Chronic exposure to cigarette smoke mimics COPD
- **Method:** Expose animals (rats/mice) to smoke over weeks/months
- **Observation:** Reduction in inflammation and lung damage = drug efficacy

B. Elastase-Induced Emphysema Model

- **Principle:** Elastase destroys alveolar walls
- **Method:** Instill elastase into lungs
- **Observation:** Prevention of alveolar destruction = protective effect

C. LPS-Induced Inflammation Model

- **Principle:** Lipopolysaccharide induces strong inflammation
- **Method:** Administer LPS into airway
- **Observation:** Reduced inflammatory markers = anti-inflammatory activity

D. Pulmonary Function Tests

- **Method:** Measure lung compliance, airway resistance
- **Observation:** Improved lung function = therapeutic effect

E. Mucus Secretion Studies

- **Method:** Measure mucus production in airway tissues
- **Observation:** Reduced mucus = beneficial in COPD

In Vitro Methods

1. Airway Smooth Muscle Studies

- Measure relaxation of bronchial muscles

2. Inflammatory Marker Assays

- **Measure cytokines like:**
  - TNF-α
  - IL-6
  - IL-8

3. Protease Inhibition Assays

- COPD involves protease-antiprotease imbalance

First Hope

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

- Evaluate inhibition of destructive enzymes

Other Alternative Models

- **Cell cultures:** Lung epithelial and immune cells
- **Zebrafish models:** Used for inflammation studies

Parameters Measured

- Lung function (FEV-like parameters in animals)
- Inflammation level
- Alveolar damage
- Mucus secretion

Toxicological Considerations

- Long-term drug safety (chronic disease treatment)
- Immunosuppression risk
- Cardiovascular side effects

Anti-Allergic Drugs – Preclinical Screening Methods

- **Allergic** reactions occur when the immune system overreacts to harmless substances (allergens).
- **This leads to:**
  - Histamine release
  - Inflammation
  - Bronchoconstriction
  - Increased vascular permeability
- **Anti-allergic drugs work by:**
  - Blocking histamine receptors (antihistamines)
  - Stabilizing mast cells
  - Reducing immune response
- Preclinical screening focuses on identifying drugs that prevent or reduce allergic reactions.

Objectives of Screening

- To evaluate anti-histaminic activity
- To assess mast cell stabilization

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.