Metabolite Identification

Metabolite Identification

Basic Idea


- Metabolite identification (Met-ID) is the process of detecting, characterizing, and determining the structure of metabolites formed after the administration of a drug.
- Metabolites are chemical compounds produced when a drug undergoes biotransformation in the body.
- Metabolite identification is an essential part of drug discovery and development because metabolites can contribute to therapeutic effects, toxicity, or drug interactions.
- The liver is the major organ involved in drug metabolism, although metabolism can also occur in the intestine, kidneys, lungs, and other tissues.

Objectives of Metabolite Identification

- To determine the metabolic pathways of a drug.
- To identify active and inactive metabolites.
- To detect toxic or reactive metabolites.
- To understand the safety profile of a drug.
- To support pharmacokinetic and toxicological studies.
- To predict possible drug-drug interactions.

First Hope

Types of Drug Metabolism



1. Phase I Reactions

- Phase I reactions involve oxidation, reduction, or hydrolysis.
- These reactions introduce or expose functional groups in the drug molecule.
- Cytochrome P450 enzymes play a major role in Phase I metabolism.

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

2. Phase II Reactions

- Phase II reactions involve conjugation with endogenous compounds such as glucuronic acid, sulfate, or glutathione.
- These reactions usually increase water solubility and facilitate excretion.

Analytical Techniques Used in Metabolite Identification

- Liquid chromatography-mass spectrometry (LC-MS).
- Liquid chromatography-tandem mass spectrometry (LC-MS/MS).
- High-resolution mass spectrometry (HRMS).
- Nuclear magnetic resonance (NMR) spectroscopy.
- Gas chromatography-mass spectrometry (GC-MS).

Importance of Metabolite Identification

- Metabolite identification improves the understanding of drug disposition.
- Metabolite identification supports the development of safer and more effective medicines.
- Metabolite identification helps regulatory agencies evaluate drug safety.

In Vitro and In Vivo Approaches

Basic Idea

- Metabolite identification studies can be performed using in vitro systems or in vivo studies.
- Both approaches provide important information about the metabolism and disposition of drugs.

In Vitro Approaches

Basic Idea

- In vitro studies are conducted outside the living organism under controlled laboratory conditions.
- These studies are widely used during early drug development.

Common In Vitro Systems

- Liver microsomes.
- Hepatocytes.
- Liver S9 fractions.
- Recombinant enzymes.
- Cell culture models.

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

Advantages

- In vitro studies are rapid and cost-effective.
- In vitro studies require smaller amounts of test compounds.
- In vitro studies reduce the use of experimental animals.

Limitations

- In vitro studies do not completely represent whole-body physiology.
- In vitro studies may not accurately predict all metabolic pathways.

In Vivo Approaches

Basic Idea

- In vivo studies are performed in living animals or humans.
- These studies provide comprehensive information about drug absorption, distribution, metabolism, and excretion.

Common Biological Samples

- Plasma.
- Urine.
- Feces.
- Bile.
- Tissue samples.

First Hope

Advantages

- In vivo studies provide realistic metabolic profiles.
- In vivo studies evaluate the combined effects of metabolism and transport processes.

Limitations

- In vivo studies are expensive and time-consuming.
- Ethical considerations are associated with animal and human studies.

Protocols and Sample Preparation

Basic Idea

- Proper protocols and sample preparation are essential for accurate metabolite identification.
- Sample preparation removes interfering substances and improves the detection of metabolites.

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

General Procedure

1. Selection of Study System

- An appropriate in vitro or in vivo model is selected according to the study objectives.

2. Drug Administration or Incubation

- The drug is administered to the study system under controlled conditions.

3. Sample Collection

- Biological samples are collected at predetermined time intervals.
- Sample collection time should cover the formation and elimination of metabolites.

4. Sample Storage

- Samples are stored under suitable conditions to prevent degradation.
- Low temperatures are commonly used to maintain stability.

5. Sample Preparation

- Proteins and interfering substances are removed before analysis.

Common Sample Preparation Techniques

A. Protein Precipitation

- Organic solvents are used to precipitate proteins from biological samples.

B. Liquid-Liquid Extraction

- Metabolites are separated based on differences in solubility between two immiscible liquids.

C. Solid Phase Extraction

- Metabolites are selectively retained on a solid sorbent and then eluted.

Sample Analysis

- Prepared samples are analyzed using techniques such as LC-MS, LC-MS/MS, or HRMS.
- Data analysis software is used to identify and characterize metabolites.

Microsomal Approaches: Rat Liver Microsomes (RLM) and Human Liver Microsomes (HLM) in Metabolite Identification

Basic Idea

- Liver microsomes are subcellular fractions obtained from the endoplasmic reticulum of liver cells.

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

- Microsomes contain important drug-metabolizing enzymes, especially cytochrome P450 enzymes and uridine diphosphate glucuronosyltransferases.
- Microsomal studies are widely used for screening metabolic pathways and predicting drug interactions.

Principle

- The drug is incubated with liver microsomes in the presence of required cofactors.
- Enzymes present in the microsomes convert the drug into metabolites.
- The generated metabolites are analyzed using analytical techniques such as LC-MS/MS.

Components of Microsomal Incubation

- Liver microsomes.
- Drug substrate.
- Buffer solution.
- Cofactors such as nicotinamide adenine dinucleotide phosphate (NADPH).

Rat Liver Microsomes (RLM)

Basic Idea

- Rat liver microsomes are prepared from rat liver tissue.
- RLM studies are commonly used during early drug development.

Applications

- Screening metabolic stability.
- Identifying major metabolites.
- Predicting potential toxicity.

Limitations

- Metabolic pathways in rats may differ from those in humans.
- Species differences may limit the prediction of human metabolism.

Human Liver Microsomes (HLM)

Basic Idea

- Human liver microsomes are prepared from human liver tissue.
- HLM studies provide information that is more relevant to human drug metabolism.

Applications

- Identification of human-specific metabolites.
- Determination of cytochrome P450 enzyme involvement.

First Hope

Copyright by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.