


Pharmaceutical incompatibilities

Pharmaceutical Incompatibilities

Definition

- Pharmaceutical incompatibility refers to undesirable interactions between drugs or between a drug and its excipients (non-active ingredients) that can affect the safety, efficacy, or stability of a pharmaceutical product.
- These incompatibilities can occur during the formulation, storage, or administration of drugs, leading to changes in physical appearance, chemical stability, or therapeutic effect.

Classification of Pharmaceutical Incompatibilities



- Pharmaceutical incompatibilities can be classified into three main types:

1. Physical Incompatibility

- **Definition:** Occurs when physical changes in appearance or properties (such as solubility, color, or texture) are observed without any chemical reaction between the components.
- **Examples:**
 - **Precipitation:** When two solutions are mixed, and an insoluble precipitate forms.
 - **Immiscibility:** When two liquids do not mix (e.g., oil and water emulsions).
 - **Liquefaction of powders:** Certain powders may liquefy upon mixing due to moisture absorption (eutectic mixtures).
- **Common Causes:** Changes in solubility, particle size, pH, or mixing of incompatible solvents or excipients.

2. Chemical Incompatibility

- **Definition:** Occurs due to a chemical reaction between the drug and other ingredients, resulting in degradation, loss of efficacy, or the formation of toxic by-products.
- **Examples:**
 - **Oxidation:** Drug degradation due to exposure to oxygen (e.g., ascorbic acid).

- **Hydrolysis:** Breakdown of drugs due to reaction with water (e.g., esters or amides).
- **Complexation:** Drug interaction leading to the formation of complexes that reduce drug availability (e.g., tetracyclines with calcium or magnesium).
- **pH-dependent degradation:** Drugs sensitive to pH changes, like aspirin, can degrade into inactive forms.
- **Common Causes:** Exposure to light, heat, moisture, pH changes, or reaction with excipients.

3. Therapeutic Incompatibility

- **Definition:** Occurs when two or more drugs administered together produce undesirable pharmacological effects or reduce each other's efficacy.
- **Examples:**
 - **Synergistic effects:** Combined drugs may lead to an exaggerated response, resulting in toxicity (e.g., alcohol and sedatives).
 - **Antagonistic effects:** One drug may reduce or negate the effect of another (e.g., antacids reducing the absorption of antibiotics).
 - **Altered absorption, distribution, metabolism, or excretion:** One drug may interfere with the pharmacokinetics of another (e.g., warfarin and certain antibiotics).
- **Common Causes:** Drug interactions due to pharmacodynamic or pharmacokinetic interference, administration route, or timing of drug intake.

Physical incompatibilities

- Physical incompatibilities involve changes in the appearance, texture, or stability of pharmaceutical formulations when substances with different physical properties are mixed.
- These incompatibilities can affect **product quality, efficacy, and safety**.

Examples of Physical Incompatibilities:

1. Immiscibility

- Occurs when two liquids do not mix to form a homogeneous solution, leading to phase separation.
- **Example:** Mixing castor oil with water results in two immiscible layers.

2. Insolubility

- Refers to a substance's inability to dissolve in a particular solvent, causing particle formation or sedimentation.
- **Example:** Barium sulfate does not dissolve in water, forming an insoluble suspension.

3. Precipitation

- The formation of solid particles from a solution, usually due to a chemical reaction or exceeding solubility limits.
- **Example:** Mixing silver nitrate with sodium chloride results in the precipitation of silver chloride.

4. Liquefaction

- Occurs when a solid substance becomes liquid due to temperature changes or interactions with other substances.
- **Example:** Camphor and menthol form a eutectic mixture, causing liquefaction.

Chemical incompatibilities

- Chemical incompatibilities involve **undesirable reactions** between substances in a pharmaceutical formulation, leading to degradation, formation of new compounds, or changes in stability, appearance, or efficacy.

Examples of Chemical Incompatibilities:

1. Alkaloid Incompatibilities

- Alkaloids can form insoluble salts when mixed with acids.
- **Example:** Morphine forms an insoluble precipitate when mixed with tannic acid.

2. Soluble Salicylate Incompatibilities

- Salicylates like aspirin can react with other substances.
- **Example:** Aspirin mixed with iron(III) ions forms a purple precipitate of ferric salicylate.

3. Soluble Iodide Incompatibilities

- Iodides can react with other compounds, forming insoluble precipitates or causing oxidation.
- **Example:** Potassium iodide mixed with lead(II) nitrate forms an insoluble yellow precipitate of lead(II) iodide.

4. Incompatibilities Causing CO₂ Evolution

- Some reactions produce carbon dioxide, affecting product stability.
- **Example:** Sodium bicarbonate and citric acid in effervescent tablets react in water, releasing CO₂ and dissolving the tablet.

5. Miscellaneous Incompatibilities

- Other reactions can lead to drug inactivation or degradation.
- **Example:** Penicillin mixed with aminoglycosides (e.g., gentamicin) leads to the inactivation of both drugs.

Therapeutic incompatibilities

- Therapeutic incompatibilities occur when two or more drugs interact, leading to **altered effects, reduced efficacy, increased toxicity, or unintended outcomes**.
- These interactions can affect **drug absorption, distribution, metabolism, or excretion (ADME)**.

Types and Causes of Therapeutic Incompatibilities:

1. Antagonism

- **Definition:** When the effect of two drugs is less than the sum of their individual effects, reducing efficacy.
- **Example:** A beta-blocker (e.g., propranolol) reduces the effect of a beta-agonist (e.g., albuterol) due to opposing actions on beta-adrenergic receptors.

2. Synergism

- **Definition:** When the combined effect of two drugs is greater than their individual effects, possibly increasing toxicity.
- **Example:** Combining a sedative (e.g., diazepam) with an opioid (e.g., morphine) can lead to excessive CNS depression and respiratory failure.

3. Altered Absorption

- **Definition:** One drug interferes with the absorption of another, reducing efficacy.
- **Example:** Tetracycline's absorption is reduced when taken with calcium or iron, as they form chelate complexes.

4. Altered Distribution

- **Definition:** One drug alters the distribution of another, affecting its concentration and activity.
- **Example:** Phenytoin displaces warfarin from protein-binding sites, increasing free warfarin and the risk of bleeding.

5. Altered Metabolism

- **Definition:** One drug affects the metabolism of another, often through enzyme induction or inhibition.
- **Example:** Rifampin induces CYP450 enzymes, increasing the metabolism of oral contraceptives, reducing their effectiveness.

6. Altered Excretion

- **Definition:** One drug changes the excretion rate of another, affecting its duration and potency.



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.