


Study of the Official Compounds

Chlorbutol (Chlorobutanol)

Chemical Structure & Formula

- **IUPAC Name:** 1,1,1-Trichloro-2-methyl-2-propanol
- **Chemical Formula:** C₄H₇Cl₃O
- **Molecular Weight:** 177.46 g/mol
- **Structure:**
 - Central carbon is bonded to:
 - Three chlorine atoms
 - One hydroxyl group (–OH)
 - One methyl group (–CH₃)
- It is a tertiary alcohol and a chlorinated derivative of butanol.



Preparation

- Chlorbutol is synthesized by the reaction of chloroform with acetone in the presence of potassium or sodium hydroxide (a base).

Reaction:



Steps:

1. Reactants:

- Chloroform (CHCl₃)
- Acetone (CH₃COCH₃)
- Sodium hydroxide (NaOH)

2. Reaction Conditions:

- Low temperature (below 20°C)
- Stirring and cooling

- Slow addition of NaOH to control the exothermic nature of the reaction

3. Mechanism:

- Base catalyzes haloform reaction between chloroform and acetone
- Trichloromethyl carbanion attacks acetone
- Hydroxylation yields chlorbutol

Test for Purity

- **Melting Point:** Pure chlorbutol has a melting point of 95–97°C.
- **Infrared spectroscopy (IR):** Peaks corresponding to –OH group ($\sim 3300\text{ cm}^{-1}$) and C–Cl stretches.
- **Thin Layer Chromatography (TLC):** Single spot indicates purity.
- **Loss on Drying:** Should be minimal to ensure absence of moisture.
- **Color of solution in alcohol:** Should remain clear without turbidity.

Assay

- **Method:** Non-aqueous titration using perchloric acid in glacial acetic acid.
- **Procedure:**
 1. Dissolve a known quantity in glacial acetic acid.
 2. Titrate with 0.1 N perchloric acid.
 3. Use crystal violet as an indicator (end-point: color change).
- **Calculation:** Based on volume used and normality.
- **Alternative:** UV/IR spectroscopy for precise quantitative analysis.

Medicinal Uses

- **Preservative:** In eye drops, ear drops, and injectable solutions (low toxicity at low concentrations).
- **Sedative & Hypnotic:** Has mild CNS depressant properties.
- **Antibacterial/Antifungal:** Used topically or in formulations to maintain sterility.


Dimercaprol (British Anti-Lewisite, BAL)

Chemical Structure & Formula

- **IUPAC Name:** 2,3-Dimercaptopropanol
- **Chemical Formula:** $\text{C}_3\text{H}_8\text{OS}_2$
- **Molecular Weight:** 124.22 g/mol

- **Structure:**

- A propane backbone
- Two adjacent thiol (–SH) groups at carbon 2 and 3
- One hydroxyl (–OH) group at carbon 1



Preparation

- Dimercaprol was originally developed as an antidote to arsenic-based chemical warfare agents.

Synthesis:

1. **Starting materials:** Allyl alcohol ($\text{CH}_2=\text{CH}-\text{CH}_2\text{OH}$)
 2. **Step 1:** React with elemental sulfur or hydrogen sulfide under reducing conditions.
 3. **Step 2:** Catalytic hydrogenation of allyl alcohol to introduce –SH groups.
- **Modern processes may involve:**
 - **Thioacetate intermediates** (e.g., reaction of epichlorohydrin with sodium thioacetate followed by hydrolysis).
 - **Note:** The product is unstable in presence of oxygen and must be stabilized in oil-based formulations (usually peanut oil).

Test for Purity

- **Odor:** Strong sulfurous odor due to thiol groups.
- **Melting point:** Should conform to standard limits ($\sim 23^\circ\text{C}$).
- **GC-MS:** For identification and purity.
- **TLC or HPLC:** Used for quantification and impurity profiling.

Assay

- **Method:** Iodometric titration
- **Principle:** The thiol groups in dimercaprol reduce iodine to iodide.
- **Procedure:**
 1. Dissolve sample in water and acetic acid.
 2. Add excess iodine solution.

3. Back titrate unreacted iodine with sodium thiosulfate using starch as an indicator.

Alternate Methods:

- High-Performance Liquid Chromatography (HPLC) with UV detection.
- Spectrophotometry based on reaction with Ellman's reagent (DTNB).


Medicinal Uses

- **Antidote for heavy metal poisoning:**
 - Arsenic, mercury, lead, and gold.
- **Chelating Agent:**
 - Binds to heavy metals through –SH groups forming stable, excretable complexes.
- **Usually administered:** Intramuscularly in peanut oil due to instability in aqueous media.
- **Cautions:** Narrow therapeutic index; causes nausea, hypertension, pain at injection site.

Glyceryl Trinitrate (Nitroglycerin)


Chemical Structure & Formula

- **IUPAC Name:** 1,2,3-Propanetriol trinitrate
- **Chemical Formula:** C₃H₅N₃O₉
- **Molecular Weight:** 227.09 g/mol
- **Structure:**
 - Glycerol molecule where all three –OH groups are esterified with nitric acid (–ONO₂).
 - Three nitrate esters on a propane backbone.
- **Highly explosive** due to nitrate groups; also a powerful vasodilator.



Preparation

- **Process:** Nitration of glycerol using a nitrating mixture (Nitric acid + Sulfuric acid)
- **Reaction:**



Steps:

1. **Cooling:** Reaction must be done at 0–5°C to control exothermic reaction.
2. Add glycerol slowly into the nitrating mixture (not vice versa).
3. Stirring and temperature control are critical to avoid explosion.
4. After reaction, mixture is poured into cold water to precipitate product.
5. Product is washed and stored in alcohol or diluted for medical use.

Test for Purity

- **Appearance:** Colorless or pale yellow liquid.
- **IR Spectroscopy:** Strong absorption bands for nitrate esters ($\sim 1630\text{--}1650\text{ cm}^{-1}$).
- **Stability Tests:** As it is volatile and sensitive, tested under stress for decomposition.
- **Limit tests:** For acidic impurities and residual sulfuric/nitric acid.

Assay

- **UV Spectrophotometry:** Nitroglycerin absorbs UV light (typically around 210–230 nm).
- **Chromatographic Methods:** HPLC is the preferred assay method.
 - Mobile Phase: Acetonitrile/water
 - Detection: UV
- **Standardization:** Compared with reference standards.

Alternative:

- Titration with sodium hydroxide after alkaline hydrolysis (liberates nitric acid equivalents).

Medicinal Uses

- **Anti-anginal agent:** Relieves chest pain by dilating coronary arteries.
- **Mechanism:**
 - Converts to nitric oxide (NO) in the body.
 - NO activates guanylate cyclase → increases cGMP → smooth muscle relaxation.
- **Dosage Forms:**
 - Sublingual tablets (fast action)
 - Ointments
 - Patches
 - Sprays
- **Used in:** Angina pectoris, heart failure, and hypertension emergencies.
- **Caution:** May cause headache, hypotension, and tolerance on prolonged use.



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.