


Diseases of Immunity

Types of Shock


What is shock?

- Shock is a life-threatening medical condition where **tissue perfusion is inadequate**, leading to **cellular hypoxia** and **organ dysfunction**.
- It represents a state of circulatory failure that impairs oxygen delivery and nutrient supply.



Types of Shock

Shock is classified based on **underlying mechanisms**, each with distinct causes and pathophysiological features:



A. Hypovolemic Shock

- **Definition:** Caused by **reduced intravascular volume**, leading to decreased preload and cardiac output.
- **Mechanism:** Decreased preload → reduced cardiac output → inadequate tissue perfusion.
- **Causes:**
 - Hemorrhage (trauma, GI bleeding, ruptured aneurysm)

- Dehydration (vomiting, diarrhea, burns)
- Third spacing (ascites, pancreatitis)

➤ **Key Features:**

- Low BP, tachycardia, cold clammy skin
- Narrow pulse pressure
- Low central venous pressure (CVP)

B. Cardiogenic Shock

- **Definition:** Failure of the heart to pump effectively despite adequate volume.
- **Mechanism:** Impaired contractility → decreased cardiac output despite normal or increased blood volume.
- **Causes:**
 - Myocardial infarction (most common)
 - Arrhythmias
 - Cardiomyopathy
 - Valve diseases
- **Key Features:**
 - Hypotension, pulmonary edema (crackles), raised JVP
 - Low cardiac output
 - High filling pressures (PCWP, CVP)

C. Distributive Shock

- **Definition:** Abnormal distribution of blood due to vasodilation.
- **Mechanism:** Blood vessels dilate, leading to relative hypovolemia even with normal/increased blood volume.
- **Subtypes:**
 - **Septic shock:** due to infection (most common distributive type)
 - **Anaphylactic shock:** IgE-mediated hypersensitivity
 - **Neurogenic shock:** spinal cord injury causing sympathetic loss
- **Key Features:**
 - **Warm extremities** (early septic shock)
 - Low systemic vascular resistance (SVR)
 - High cardiac output (initially in sepsis)

D. Obstructive Shock

- **Definition:** Physical obstruction to circulation.
- **Mechanism:** Reduced cardiac output due to mechanical blockage.
- **Causes:**
 - Pulmonary embolism
 - Cardiac tamponade
 - Tension pneumothorax
 - Constrictive pericarditis
- **Key Features:**
 - Hypotension with signs of obstruction
 - Raised JVP (tamponade/PE)
 - Pulsus paradoxus (in tamponade)

Mechanisms of Shock

- Shock leads to tissue hypoxia through multiple interconnected mechanisms that can escalate to **Multiple Organ Dysfunction Syndrome (MODS)** if not corrected.
- These mechanisms can be grouped into pathological processes and compensatory responses:

Pathophysiological Mechanisms

1. Inadequate Perfusion

- ↓ Oxygen and nutrient delivery to tissues impairs cellular metabolism.

2. Anaerobic Metabolism

- Oxygen deficiency forces cells into anaerobic glycolysis → **lactic acid accumulation** → **metabolic acidosis**.

3. Cell Membrane Dysfunction

- Failure of ATP-dependent ion pumps
- Sodium and water influx → **cellular edema**
- Calcium influx → **cell death**

4. Systemic Inflammatory Response (*particularly prominent in septic shock*)

- Cytokine release
- Endothelial dysfunction
- Capillary leak and vasodilation
- Coagulopathy (may contribute to DIC)

Compensatory Mechanisms

- These short-term responses aim to maintain perfusion and blood pressure:

A. Sympathetic Nervous System Activation

- → Vasoconstriction, tachycardia, ↑ cardiac output.

B. Renin-Angiotensin-Aldosterone System (RAAS) Activation

- → Salt and water retention to expand intravascular volume.

C. Antidiuretic Hormone (ADH) Release


- → Promotes water reabsorption in the kidneys → fluid retention.

Progression to MODS (Multiple Organ Dysfunction Syndrome)

- If shock remains uncorrected:
 - Initial compensatory mechanisms fail
 - Hypoperfusion worsens
 - Inflammatory and cellular injury becomes systemic
 - **MODS develops**, involving failure of vital organs such as lungs, kidneys, liver, and heart.

Stages of Shock

Shock progresses through several stages:



A. Initial Stage (Pre-shock or Early Shock)

Physiological Changes:

- Reduced tissue perfusion
- No visible clinical symptoms



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.