

Patient medication adherence

Definition of Patient Medication Adherence

What is Medication Adherence?

- Medication adherence refers to the degree to which a patient correctly follows medical advice and instructions related to prescribed medications. This includes:
 - Taking medications at the right time
 - In the correct dose
 - For the full duration prescribed
 - In the correct way (e.g., with food, on an empty stomach, etc.)
- According to the **World Health Organization (WHO)**, adherence is defined as:
 - *"The extent to which a person's behavior—taking medication, following a diet, and/or executing lifestyle changes—corresponds with agreed recommendations from a healthcare provider."*

Distinction Between Adherence and Compliance

- **Compliance** implies a more passive role of the patient, simply following the doctor's instructions.
- **Adherence**, on the other hand, emphasizes a partnership between the patient and the healthcare provider, where the patient is actively involved in the decision-making process.

Types of Medication Adherence Behavior:

1. **Adherent** – Patient takes medication exactly as prescribed.
2. **Non-adherent** – Patient does not follow the prescribed regimen.
 - **Intentional non-adherence:** Deliberate choice (e.g., due to side effects, beliefs).
 - **Unintentional non-adherence:** Due to forgetfulness, misunderstanding, etc.

Types of Non-Adherence

- There are two major types:
 1. **Intentional non-adherence:** The patient consciously chooses not to follow the treatment plan. Reasons may include beliefs about the medicine, fear of side effects, or perceived lack of benefit.
 2. **Unintentional non-adherence:** Occurs when the patient wants to follow the treatment plan but fails due to external factors such as forgetfulness, misunderstanding instructions, or financial constraints.

Factors Affecting Medication Adherence

- Medication adherence is influenced by a complex set of factors.

WHO categorizes these into five key dimensions:

A. Socioeconomic Factors

- These include the broader social and economic circumstances in which a patient lives.
 - **Low income or poverty:** Medication may be unaffordable.
 - **Lack of health insurance** or coverage limitations.
 - **Low educational level:** Difficulty understanding prescription instructions.
 - **Lack of social support:** No family or caregiver to assist.
 - **Cultural beliefs and practices:** May influence a patient's views on Western medicine or certain drug therapies.

B. Health System/Healthcare Team-Related Factors

- These involve how the healthcare system and providers interact with the patient.
 - **Poor communication between patient and provider:** If patients don't understand their diagnosis or treatment plan, they may not adhere properly.
 - Long wait times or inaccessible pharmacy services.
 - **Lack of continuity of care:** Seeing different doctors may result in confusion or conflicting advice.
 - **Limited consultation time:** Pharmacists or doctors may not have enough time to explain the importance of adherence.

C. Condition-Related Factors

- These relate to the patient's illness or health condition.
 - **Chronic diseases** (e.g., hypertension, diabetes): Long-term therapy may lead to fatigue or forgetfulness.
 - **Asymptomatic conditions:** Patients may not feel sick (e.g., high blood pressure) and stop taking medication.
 - **Mental health conditions:** Depression or cognitive disorders can interfere with a patient's ability to remember or manage medication.

D. Therapy-Related Factors

- These include the complexity and characteristics of the prescribed therapy.
 - **Complex regimens:** Multiple medications or doses throughout the day can be difficult to manage.
 - **Side effects:** Fear or experience of adverse drug reactions may lead to discontinuation.

- **Duration of treatment:** Longer courses can lead to declining adherence over time.
- **Route of administration:** Injectable or inhalable medications may be more difficult for patients to use correctly.

E. Patient-Related Factors

- These are individual patient characteristics and behaviors.
 - Forgetfulness or busy lifestyle.
 - Lack of motivation or understanding of the importance of adherence.
 - Fear of dependence on medication.
 - Misconceptions about disease and treatment.
 - **Personal beliefs and preferences:** Some may prefer herbal remedies or fear pharmaceuticals.

Formula for Medication Adherence

Medication Adherence Rate (%)

$$\text{Adherence (\%)} = \left(\frac{\text{Number of doses taken}}{\text{Number of doses prescribed}} \right) \times 100$$

Explanation:

- **Number of doses taken:** The actual number of doses the patient consumed over a given period.
- **Number of doses prescribed:** The total number of doses the healthcare provider prescribed for that same period.
- **Adherence (%):** Expressed as a percentage, where 100% indicates perfect adherence.

Example:

- If a patient was prescribed **2 doses per day for 30 days**, the total number of prescribed doses is:

$$2 \times 30 = 60 \text{ doses}$$

- If the patient actually took **54 doses**:

$$\text{Adherence} = \left(\frac{54}{60} \right) \times 100 = 90\%$$

- This means the patient has a **90% adherence rate**.

Role of the Pharmacist in Improving Medication Adherence

- Community pharmacists play a central and direct role in improving patient medication adherence. As one of the most accessible healthcare professionals, pharmacists are in a

unique position to interact regularly with patients, provide education, and monitor medication use.

- Pharmacists can intervene at multiple stages of a patient’s medication journey, from dispensing to follow-up, to ensure adherence is optimized.

Patient Education and Counseling

One of the most important responsibilities of a community pharmacist is to **educate and counsel** patients effectively.

A. Clear Communication

- Use simple, non-technical language when explaining how to take medications.
- Confirm the patient understands the **dose, frequency, route, and duration**.
- Use the **"teach-back method"**: Ask patients to repeat instructions in their own words to ensure comprehension.

B. Explain Importance of Adherence

- Emphasize how regular use of medication controls the disease and prevents complications.
- Help patients understand that skipping doses or stopping early may worsen the condition.

C. Address Misconceptions

- Correct common myths about side effects or dependency.
- Reassure patients about the safety and effectiveness of their prescribed therapy.

Simplifying the Medication Regimen

- Pharmacists can work with prescribers to recommend **simplified regimens**, which are easier for patients to follow.
 - **Reduce pill burden**: Suggest combination medications where available.
 - **Once-daily dosing**: Advocate for medications that can be taken less frequently.
 - **Fixed-dose combinations (FDCs)**: Help patients manage multiple conditions with fewer tablets.

Identifying and Addressing Barriers to Adherence

Pharmacists must proactively identify and address barriers that may lead to non-adherence.

Financial Constraints

- Recommend generic alternatives or more affordable brands.
- Guide patients towards **government schemes** or **insurance programs** that reduce out-of-pocket costs.

Side Effects

- Counsel patients on expected side effects and how to manage them.



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.