


Antimalarials

Etiology (Causes) of Malaria

- Malaria is caused by **Plasmodium** parasites, transmitted by female **Anopheles mosquitoes**.
- The main species causing malaria in humans are:
 1. **Plasmodium falciparum**: Most severe, can cause cerebral malaria.
 2. **Plasmodium vivax**: Causes relapses due to liver hypnozoites.
 3. **Plasmodium ovale**: Similar to P. vivax, but less common.
 4. **Plasmodium malariae**: Can cause chronic infection.
 5. **Plasmodium knowlesi**: Zoonotic malaria, found in Southeast Asia.

Transmission & Life Cycle



1. Mosquito Stage

- The mosquito injects **sporozoites** into the human bloodstream during a bite.

2. Liver Stage (Pre-Erythrocytic Stage)

- Sporozoites invade liver cells, multiply, and form **merozoites**.

3. Blood Stage (Erythrocytic Cycle)

- Merozoites infect red blood cells (RBCs), multiply, rupture RBCs, and cause fever.

4. Gametocyte Formation

- Some parasites develop into **gametocytes**, which are taken up by a mosquito.

5. Mosquito Cycle

- Gametocytes mature into **sporozoites**, completing the cycle.

Symptoms of Malaria


- High fever, chills, sweating (classic **tertian or quartan fever cycles**)
- Headache, muscle pain, fatigue
- Severe malaria (*P. falciparum*) → Organ failure, coma, death

Prevention & Treatment

- **Prevention:** Mosquito control (nets, repellents), prophylactic drugs (chloroquine, doxycycline).
- **Treatment:** Antimalarial drugs (artemisinin-based combination therapies, chloroquine, mefloquine).

Classification of Antimalarial Drugs

- Antimalarial drugs are classified based on their chemical structure and mechanism of action. The main classes include:



1. Quinolines

- These are synthetic or natural compounds that interfere with the parasite's ability to digest hemoglobin.
- **Examples:** Quinine sulphate, Chloroquine, Amodiaquine, Primaquine phosphate, Pamaquine, Quinacrine hydrochloride, Mefloquine.

Structural Feature	SAR Insight
Quinoline nucleus (1- or 4-substituted)	Essential for antimalarial activity. Most active compounds are 4- or 8-aminoquinolines.
Position 4 (4-amino group)	Common in chloroquine and amodiaquine ; important for blood schizonticidal activity.
Position 8 (8-amino group)	Seen in primaquine , pamaquine ; responsible for tissue schizonticidal and gametocidal action.
Side chain on amino group	Chain length (typically 3 C atoms) is critical. Bulky or very long chains reduce activity.

Position 6 (methoxy or other substitutions)	Methoxy or halogen (Cl, Br) often improves lipophilicity and activity (e.g., in primaquine, amodiaquine).
Electron-donating groups on ring	Increase antimalarial activity and reduce toxicity.
Electron-withdrawing groups on ring	Often reduce activity.
Basic nitrogen in side chain	Enhances accumulation in the acidic food vacuole of the parasite. Protonation helps trap the drug.
Hydroxylation or polar groups	May reduce blood stage activity but are tolerated in tissue stage drugs (e.g., primaquine).

2. Biguanides and Dihydrotriazines

- These drugs act by inhibiting dihydrofolate reductase (DHFR), which is essential for the parasite's DNA synthesis.
- **Examples:** Cycloguanil pamoate, Proguanil.

3. Miscellaneous

- This group consists of antimalarial agents with varied mechanisms, such as inhibiting mitochondrial electron transport, disrupting heme detoxification, or acting on parasite protein synthesis.
- **Examples:** Pyrimethamine, Artesunate, Artemether, Atovaquone.

Brand Names Of Important Marketed Products And Their Side Effects.

These drugs are used for the **treatment and prevention of malaria**, caused by *Plasmodium* species.

Drug Name	Brand Name(s)	Formulation	Use
Chloroquine	Lariago, Aralen	Tablet, Injection	Treatment/prophylaxis (sensitive strains)
Hydroxychloroquine	Plaquenil, HCQS	Tablet	Malaria, rheumatoid arthritis, lupus
Artemether + Lumefantrine	Coartem, Lumerax	Tablet	Uncomplicated falciparum malaria
Artesunate	Falcigo, Malinate	Tablet, Injection	Severe/complicated malaria
Mefloquine	Lariam, Mefliam	Tablet	Prophylaxis and treatment
Quinine	Quinimax, Quinidine	Injection, Tablet	Severe malaria


Primaquine	Primaquine	Tablet	Radical cure (<i>P. vivax</i> , <i>P. ovale</i>)
Atovaquone + Proguanil	Malarone	Tablet	Prophylaxis and treatment

Various Quinolines drugs

1. Quinine Sulfate

Chemical Structure / Formula

- **Formula:** C₂₀H₂₄N₂O₂ · H₂SO₄
- **Structure:** A **quinoline nucleus** with an attached quinuclidine ring.



Mechanism of Action

- Interferes with **heme polymerization** in the malaria parasite, leading to toxic heme accumulation and parasite death.
- Also **interferes with DNA synthesis** in Plasmodium spp.

Uses

- **Severe malaria** (especially against chloroquine-resistant *P. falciparum*).
- **Nocturnal leg cramps** (off-label).

Side Effects

- **Cinchonism** (tinnitus, dizziness, headache, nausea).
- **Hypoglycemia** (stimulates insulin release).
- **Hemolysis in G6PD deficiency** (Blackwater fever).
- **QT prolongation** (cardiac toxicity).

2. Chloroquine*

Chemical Structure / Formula



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.