Redox titrations

Introduction to Redox Titrations

- Redox titrations, a form of volumetric analysis, are widely used in pharmaceutical analysis to determine the concentration of substances via oxidation-reduction reactions.
- These titrations play a key role in ensuring the quality, safety, and potency of drug substances and products.

Principles of Redox Titration

1. Redox Reaction

- Redox titration is based on a reaction between an oxidizing agent and a reducing agent.
- During titration, the titrant is added to the analyte until the reaction reaches completion, often indicated by a color change or electrode potential shift.

2. Indicators

- Indicators are frequently used to signal the endpoint of the titration, where the amount of titrant added is stoichiometrically equivalent to the analyte.

3. Electrochemical Methods

- In certain cases, particularly with complex systems or low concentrations, electrochemical methods like potentiometry are used to detect the endpoint by measuring the potential difference between electrodes.

Applications in Pharmaceutical Analysis

1. Determination of Active Pharmaceutical Ingredients (APIs):

- Redox titrations are used to quantify APIs, ensuring correct dosage and drug efficacy.

2. Impurity Profiling:

- These titrations help identify impurities that may be oxidizable or reducible, critical for drug purity and safety.

3. Stability Testing:

- Redox titrations assess drug stability by monitoring degradation, which often involves redox reactions over time.

4. Formulation Development:

- During drug development, redox titration is essential for optimizing formulations involving redox-active components.

Advantages in Pharmaceutical Analysis

1. Accuracy and Precision:

Copyright © 2025 by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

- Redox titrations offer highly accurate and precise measurements, essential for quality control and regulatory compliance.

2. **Versatility:**

- They can be applied to a broad range of substances, including those challenging to analyze by other techniques.

3. **Sensitivity:**

- These titrations can detect minute changes in the redox state, making them highly sensitive and reliable.

Concept of Oxidation

- Oxidation is a process in which a chemical species **loses one or more electrons**, leading to an **increase in its oxidation state**.
- In simpler terms, the species becomes more positively charged.
- Oxidation is often associated with the **addition of oxygen** to a substance or the **removal of hydrogen** from it.
- However, the key factor that defines oxidation is the **loss of electrons**.

Example:

- Consider the reaction between magnesium metal (Mg) and oxygen gas (O_2) to form magnesium oxide (MgO).

$$2\,Mg\,(s) + O_2\,(g) \longrightarrow 2\,MgO\,(s)$$

- In this reaction, magnesium loses two electrons to become the Mg^{2+} ion, as shown below:

$$Mg\,(s) \longrightarrow Mg^{2+}(s) + 2e^-$$

- Magnesium has been oxidized, as it has lost electrons, and its oxidation state has increased from 0 to +2.

Oxidizing Agents

- An **oxidizing agent** is a substance that gains electrons in a redox reaction, causing the oxidation of another species.
- By accepting electrons from another species, it enables that species to lose electrons, leading to its oxidation.
- As a result, the oxidizing agent itself gets reduced in the process.
- Oxidizing agents typically have a high affinity for electrons and can be identified by their ability to oxidize other substances.

Example:

Copyright © 2025 by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

- In the reaction between hydrogen gas (H_2) and oxygen gas (O_2) to form water (H_2O):

$$2\,H_2(g) + O_2(g) \longrightarrow 2\,H_2O(l)$$

- Oxygen (O_2) is the **oxidizing agent**, as it gains electrons from hydrogen (H_2) during the reaction, leading to the reduction of oxygen and the oxidation of hydrogen.

Concept of Reduction

- Reduction is a process in which a chemical species **gains one or more electrons**, leading to a **decrease in its oxidation state**.
- In other words, the species becomes **less positively charged or more negatively charged**.
- Reduction is often associated with the **addition of hydrogen** to a substance or the **removal of oxygen** from it.
- Fundamentally, however, reduction is defined by the **gain of electrons**.

Example

- In the reaction between magnesium metal and oxygen gas, oxygen is reduced to form oxide ions (O^{2-}):

$$\frac{1}{2}O_2\,(g) + 4e^- \longrightarrow 2\,O^{2-}\,(s)$$

- Oxygen has **gained two electrons** in this process, and its oxidation state has decreased from 0 to –2.

Reducing Agents

- A **reducing agent** is a substance that **donates electrons** in a redox reaction, causing the reduction of another species.
- By giving away electrons to another species, it enables that species to gain electrons, leading to its reduction.
- As a result, the **reducing agent itself gets oxidized** in the process.
- Reducing agents typically have a **low affinity for electrons** and are identified by their ability to reduce other substances.

Example

- In the reaction between hydrogen gas and oxygen gas to form water:

$$2\,H_2\,(g) + O_2\,(g) \longrightarrow 2\,H_2O\,(l)$$

- Hydrogen (H_2) is the **reducing agent**, as it donates electrons to oxygen (O_2) during the reaction, leading to the **oxidation of hydrogen** and the **reduction of oxygen**.

Types of redox titrations

Copyright © 2025 by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

A. Cerimetry

Principle

- Cerimetry involves the use of **cerium(IV) salts** (e.g., cerium(IV) sulfate) as the oxidizing titrant.
- Cerium(IV) ions are reduced to cerium(III) ions when they react with reducing agents in the analyte.

Applications

- Determination of **reducing sugars**, **phenols**, and various **organic and inorganic substances**.

Example Chemical Reaction

$$2\,Ce(SO_4)_2 + C_2H_2O_4 \longrightarrow 2\,CeSO_4 + 2\,H_2SO_4 + 2\,CO_2$$

B. Iodimetry

Principle

- Iodimetry involves the **direct titration of a reducing agent with iodine (I_2)** as the oxidizing titrant.
- Iodine is reduced to iodide (I^-), while the reducing agent is oxidized.

Applications

- Used to quantify reducing agents such as **sulfites, thiosulfates, arsenites**, and organic compounds like **vitamin C**.

Example Chemical Reaction

$$2\,S_2O_3^{2-} + I_2 \longrightarrow S_4O_6^{2-} + 2\,I^-$$

C. Iodometry

Principle

- Iodometry is an **indirect redox titration** where the reducing agent reacts with **excess iodine**, producing iodide ions.
- The unreacted iodine is then titrated with **sodium thiosulfate**.

Applications

- Determination of **metal ions** (e.g., copper, antimony) and various **organic and pharmaceutical substances**.

Example Chemical Reaction

1. Copper(II) sulfate reacts with potassium iodide:

$$Cu^{2+} + 2\,I^- \longrightarrow CuI\,(s) + I_2$$

2. Liberated iodine is titrated with sodium thiosulfate:

Copyright © 2025 by Engacy Learning Private Limited. All rights reserved. Reproduction or distribution without written consent is strictly prohibited.

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.