Patient medication adherence

Definition of Patient Medication Adherence

What is Medication Adherence?

- Medication adherence refers to the degree to which a patient correctly follows medical advice and instructions related to prescribed medications. This includes:
    - Taking medications at the right time
    - In the correct dose
    - For the full duration prescribed
    - In the correct way (e.g., with food, on an empty stomach, etc.)
- According to the **World Health Organization (WHO)**, adherence is defined as:
    - *"The extent to which a person's behavior—taking medication, following a diet, and/or executing lifestyle changes—corresponds with agreed recommendations from a healthcare provider."*

Distinction Between Adherence and Compliance

- **Compliance** implies a more passive role of the patient, simply following the doctor's instructions.
- **Adherence**, on the other hand, emphasizes a partnership between the patient and the healthcare provider, where the patient is actively involved in the decision-making process.

Types of Medication Adherence Behavior:

1. **Adherent** – Patient takes medication exactly as prescribed.
2. **Non-adherent** – Patient does not follow the prescribed regimen.
    - **Intentional non-adherence**: Deliberate choice (e.g., due to side effects, beliefs).
    - **Unintentional non-adherence**: Due to forgetfulness, misunderstanding, etc.

Types of Non-Adherence

- **There are two major types:**
    1. **Intentional non-adherence**: The patient consciously chooses not to follow the treatment plan. Reasons may include beliefs about the medicine, fear of side effects, or perceived lack of benefit.
    2. **Unintentional non-adherence**: Occurs when the patient wants to follow the treatment plan but fails due to external factors such as forgetfulness, misunderstanding instructions, or financial constraints.

Factors Affecting Medication Adherence

- Medication adherence is influenced by a complex set of factors.

WHO categorizes these into five key dimensions:

A. Socioeconomic Factors

- These include the broader social and economic circumstances in which a patient lives.
  - **Low income or poverty**: Medication may be unaffordable.
  - **Lack of health insurance** or coverage limitations.
  - **Low educational level**: Difficulty understanding prescription instructions.
  - **Lack of social support**: No family or caregiver to assist.
  - **Cultural beliefs and practices**: May influence a patient's views on Western medicine or certain drug therapies.

B. Health System/Healthcare Team-Related Factors

- These involve how the healthcare system and providers interact with the patient.
  - **Poor communication between patient and provider**: If patients don't understand their diagnosis or treatment plan, they may not adhere properly.
  - Long wait times or inaccessible pharmacy services.
  - **Lack of continuity of care**: Seeing different doctors may result in confusion or conflicting advice.
  - **Limited consultation time**: Pharmacists or doctors may not have enough time to explain the importance of adherence.

C. Condition-Related Factors

- These relate to the patient's illness or health condition.
  - **Chronic diseases** (e.g., hypertension, diabetes): Long-term therapy may lead to fatigue or forgetfulness.
  - **Asymptomatic conditions**: Patients may not feel sick (e.g., high blood pressure) and stop taking medication.
  - **Mental health conditions**: Depression or cognitive disorders can interfere with a patient's ability to remember or manage medication.

D. Therapy-Related Factors

- These include the complexity and characteristics of the prescribed therapy.
  - **Complex regimens**: Multiple medications or doses throughout the day can be difficult to manage.
  - **Side effects**: Fear or experience of adverse drug reactions may lead to discontinuation.

- **Duration of treatment**: Longer courses can lead to declining adherence over time.
- **Route of administration**: Injectable or inhalable medications may be more difficult for patients to use correctly.

E. Patient-Related Factors

- These are individual patient characteristics and behaviors.
  - Forgetfulness or busy lifestyle.
  - Lack of motivation or understanding of the importance of adherence.
  - Fear of dependence on medication.
  - Misconceptions about disease and treatment.
  - **Personal beliefs and preferences**: Some may prefer herbal remedies or fear pharmaceuticals.

Formula for Medication Adherence

Medication Adherence Rate (%)

$$\text{Adherence (\%)} = \left(\frac{\text{Number of doses taken}}{\text{Number of doses prescribed}}\right) \times 100$$

Explanation:

- **Number of doses taken**: The actual number of doses the patient consumed over a given period.
- **Number of doses prescribed**: The total number of doses the healthcare provider prescribed for that same period.
- **Adherence (%)**: Expressed as a percentage, where 100% indicates perfect adherence.

Example:

- If a patient was prescribed **2 doses per day for 30 days**, the total number of prescribed doses is:

$$2 \times 30 = 60 \text{ doses}$$

- If the patient actually took **54 doses**:

$$\text{Adherence} = \left(\frac{54}{60}\right) \times 100 = 90\%$$

- This means the patient has a **90% adherence rate**.

Role of the Pharmacist in Improving Medication Adherence

- Community pharmacists play a central and direct role in improving patient medication adherence. As one of the most accessible healthcare professionals, pharmacists are in a

unique position to interact regularly with patients, provide education, and monitor medication use.

- Pharmacists can intervene at multiple stages of a patient's medication journey, from dispensing to follow-up, to ensure adherence is optimized.

Patient Education and Counseling

One of the most important responsibilities of a community pharmacist is to **educate and counsel** patients effectively.

A. Clear Communication

- Use simple, non-technical language when explaining how to take medications.
- Confirm the patient understands the **dose, frequency, route, and duration**.
- Use the **"teach-back method"**: Ask patients to repeat instructions in their own words to ensure comprehension.

B. Explain Importance of Adherence

- Emphasize how regular use of medication controls the disease and prevents complications.
- Help patients understand that skipping doses or stopping early may worsen the condition.

C. Address Misconceptions

- Correct common myths about side effects or dependency.
- Reassure patients about the safety and effectiveness of their prescribed therapy.

Simplifying the Medication Regimen

- Pharmacists can work with prescribers to recommend **simplified regimens**, which are easier for patients to follow.
  - **Reduce pill burden**: Suggest combination medications where available.
  - **Once-daily dosing**: Advocate for medications that can be taken less frequently.
  - **Fixed-dose combinations (FDCs)**: Help patients manage multiple conditions with fewer tablets.

Identifying and Addressing Barriers to Adherence

Pharmacists must proactively identify and address barriers that may lead to non-adherence.

Financial Constraints

- Recommend generic alternatives or more affordable brands.
- Guide patients towards **government schemes** or **insurance programs** that reduce out-of-pocket costs.

Side Effects

- Counsel patients on expected side effects and how to manage them.

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.