Column Chromatography

Column Chromatography


Introduction

- Column Chromatography is a classical separation technique used to isolate, identify, and purify chemical compounds.
- It is widely used in pharmaceutical analysis for the separation of complex mixtures based on the differential interactions of components with the stationary and mobile phases.
- **Column chromatography can be broadly classified into:**
    - Adsorption column chromatography
    - Partition chromatography

Adsorption Column Chromatography

Principle

- This method is based on adsorption of analytes onto a solid stationary phase (usually silica gel or alumina), and their desorption by a liquid mobile phase.
- Components are separated due to differences in their affinity for the stationary phase.



Stationary Phase

- **Adsorbent**: Silica gel (polar), Alumina (polar)
- Nature: Solid, highly porous

Mobile Phase

- **Eluent**: Organic solvents like hexane, chloroform, methanol, ethyl acetate, etc.
- Selected based on polarity and elution strength

Mechanism

- Components with stronger adsorption elute slower (higher retention)
- Components with weaker adsorption elute faster

Operational Technique of Column Chromatography

Step-by-Step Procedure

1. Column Packing

- Wet or dry packing of the adsorbent into the glass column
- Ensure no air bubbles or cracks

2. Sample Application

- Dissolve sample in a small volume of solvent and apply it carefully on top of the packed bed

3. Elution

- Mobile phase is added continuously
- Gravity or slight pressure helps flow
- Two common modes:
  - **Isocratic**: Constant composition
  - **Gradient**: Increasing elution strength

4. Fraction Collection

- Eluent is collected in test tubes or vials
- Monitored by UV, TLC, or HPLC

5. Detection and Analysis

- Fractions are analyzed to identify and isolate the desired compound

Frontal Analysis and Elution Analysis

These are two approaches to understand and model the separation process in column chromatography:

A. Frontal Analysis

- In this method, the sample is continuously fed into the column.
- Components move at different speeds and exit the column one after another.
- Each component appears as a plateau, not a peak.
- The order of elution is from least adsorbed to most adsorbed.
- Not used for analytical purposes; more for capacity and adsorption studies.
- **Use**: Determination of adsorption isotherms, capacity factor

B. Elution Analysis

- Most widely used form of chromatography.
- A finite quantity of sample is introduced.

- Components elute as discrete peaks.
- The component with the lowest affinity elutes first.
- **Use**: Analytical and preparative separation of pharmaceuticals

Factors Affecting Column Efficiency

1. Particle Size of Stationary Phase

- Smaller particle size → larger surface area → better resolution
- But may increase back pressure

2. Column Length and Diameter

- Longer columns improve resolution but increase time
- Narrower columns give sharper peaks

3. Uniformity of Packing

- Poor packing causes channeling and tailing
- Uniform packing improves efficiency

4. Flow Rate of Mobile Phase

- Too fast → incomplete separation
- Too slow → broad peaks and long analysis time

5. Nature of Stationary and Mobile Phase

- Polarity plays a crucial role
- Proper match improves selectivity and resolution

6. Temperature

- Affects viscosity and interaction strength
- Generally, higher temperature reduces retention time

Applications of Column Chromatography

Area	Application
Purification	Separation of synthetic drug mixtures
Isolation	Extraction of natural products like alkaloids, glycosides
Qualitative Analysis	Identification of compounds using elution time and behavior
Quantitative Analysis	Estimation of active ingredients or impurities
Preparative Chromatography	Large-scale purification for drug development
Stability Studies	Monitoring degradation products over time

Partition Chromatography

Principle

- In partition chromatography, separation is based on the partitioning of solutes between two liquid phases: a stationary liquid coated on a solid support and a mobile liquid or gas.

Key Features

- Stationary phase: Liquid film (e.g., water or alcohol)
- Mobile phase: Another immiscible liquid (in L-L chromatography) or a gas (in gas-liquid chromatography)
- Separation is based on relative solubilities of components in the two phases

Examples in Pharmaceutical Analysis

- Paper Chromatography (liquid-liquid)
- Gas-Liquid Chromatography (already discussed)
- Centrifugal partition chromatography (CPC)

Applications

- Isolation of polar and non-polar drug components
- Separation of closely related compounds
- Useful for heat-sensitive and labile substances

Comparison: Adsorption vs Partition Chromatography

Feature	Adsorption Chromatography	Partition Chromatography
Stationary Phase	Solid adsorbent (e.g., silica)	Liquid coated on inert solid
Mobile Phase	Liquid	Liquid or gas
Separation Basis	Adsorption strength	Solubility (partition coefficient)
Example	Column Chromatography	Paper Chromatography, GLC
Used For	Non-polar to moderately polar compounds	Polar and labile compounds

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.