Thin Layer Chromatography (TLC)

Thin Layer Chromatography (TLC)


Introduction

- Thin Layer Chromatography (TLC) is a simple, rapid, and versatile chromatographic technique used for the separation and identification of components in a mixture.
- It is particularly important in pharmaceutical analysis due to its cost-effectiveness, speed, and reliability.
- TLC is a form of planar chromatography, where the stationary phase is supported on a flat surface (usually a glass or plastic plate).

Principle of TLC

- TLC works on the principle of adsorption and partition:
  - The stationary phase (e.g., silica gel or alumina) adsorbs the components of the sample.
  - The mobile phase (a suitable solvent or mixture) moves by capillary action across the plate.
  - The separation occurs due to differences in the affinity (adsorption and solubility) of each component toward the stationary and mobile phases.
- Components with higher affinity for the stationary phase move slowly, while those with greater solubility in the mobile phase move faster.

Technique of TLC



A. Preparation of the Plate

- A thin layer (0.1–0.25 mm) of adsorbent (silica gel, alumina, cellulose) is spread on a plate of glass, aluminum, or plastic.
- Plates can be commercially available pre-coated or manually prepared.

B. Activation

- The plates are heated in an oven at 100–120°C for about 30 minutes to remove moisture.

C. Sample Application

- Small spots (1–2 µL) of the sample solution are applied with a capillary tube or microsyringe near the bottom edge of the plate (~1 cm above the base).
- Multiple samples or standards can be spotted side by side.

D. Development of the Plate

- The plate is placed in a development chamber (e.g., a glass jar with a lid) containing the mobile phase solvent.
- The chamber is saturated with solvent vapor using a filter paper liner to ensure uniform movement.
- The solvent moves up the plate by capillary action, carrying the components with it.
- Development is stopped when the solvent front reaches near the top (~75% height of the plate).

E. Visualization

- After development, the solvent front is marked and the plate is dried.
- Spots are visualized using:
  - UV light (254 or 366 nm) if compounds are UV-active
  - Spray reagents (e.g., ninhydrin for amino acids, iodine vapors, sulfuric acid charring)
  - Fluorescence under UV if fluorescent indicator is present in the stationary phase

Retention Factor (Rf Value)

- The **Rf value** is a characteristic parameter used to identify compounds.

Formula:

$$Rf = \frac{\text{Distance traveled by the solute}}{\text{Distance traveled by the solvent front}}$$

Key Points:

- Rf is always between 0 and 1
- It is unitless
- It varies with solvent system, adsorbent type, thickness of layer, temperature, and humidity

Interpretation:

- Higher Rf: Indicates less polar compounds (move faster)
- Lower Rf: Indicates more polar compounds (stronger interaction with stationary phase)

Importance in Pharma:

- Comparison of Rf values of sample and reference under the same conditions helps identify unknowns.

Applications of TLC in Pharmaceutical Analysis

TLC is used both qualitatively and semi-quantitatively in pharma labs.

A. Identification of Drugs and Excipients

- Compare Rf of sample with that of reference standard.

B. Detection of Impurities and Degradation Products

- Useful in stability testing and impurity profiling.

C. Analysis of Herbal Drugs

- For fingerprint profiling of phytoconstituents.

D. Monitoring of Reactions

- Used in organic synthesis to track reaction progress.

E. Purity Testing

- Presence of additional spots indicates impurities.

F. Qualitative Estimation

- Estimation of content by spot intensity comparison with standard.

G. Bioanalysis

- Detection of metabolites in biological fluids (rare due to limited sensitivity).

H. Preparative TLC

- Scraping off and eluting separated components for further testing.

Advantages of TLC

- Simple and cost-effective
- Fast with no complex instrumentation
- Multiple samples can be analyzed simultaneously
- Useful for thermally unstable and non-volatile compounds

Limitations of TLC

- Low sensitivity compared to HPLC or GC
- Rf values may vary with experimental conditions
- Not suitable for quantitative analysis in routine pharma QC
- Manual method prone to operator error

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.