High-Performance Thin Layer Chromatography (HPTLC)

High-Performance Thin Layer Chromatography (HPTLC)

Introduction

- High-Performance Thin Layer Chromatography (HPTLC) is an advanced and automated form of traditional Thin Layer Chromatography (TLC).
- It offers greater resolution, sensitivity, reproducibility, and speed, making it suitable for qualitative and quantitative analysis in pharmaceutical and biomedical research.
- HPTLC is widely accepted by pharmacopeias (USP, IP, BP) and regulatory authorities for:
    - Identification of active pharmaceutical ingredients (APIs)
    - Estimation of impurities and degradation products
    - Quality control of herbal formulations
- It is especially useful in multi-component analysis and routine quality control where high sample throughput is required.

Theory of HPTLC

- HPTLC is based on the same fundamental principle as TLC:
    - Partition and adsorption chromatography, where separation occurs due to differences in the affinities of components toward the stationary and mobile phases.
- **In HPTLC:**
    - The stationary phase is a high-resolution, precoated layer of fine particle silica or alumina (5–10 µm) with a uniform and dense coating.
    - The mobile phase is a solvent or solvent mixture that moves via capillary action.
    - The components of a sample partition between the two phases and separate based on their polarity, solubility, and molecular interaction.
- Because of its controlled environment and automation, HPTLC allows precise migration, well-defined Rf values, and quantitative data output.

Instrumentation of HPTLC

The HPTLC system consists of modular, computer-controlled units, which offer high precision and reproducibility.

1. Sample Applicator (Automatic Spotter)

- Precisely applies the sample as narrow bands or spots using a micro-syringe or capillary system.
- Ensures uniformity, which is critical for quantitative analysis.
- Multiple samples can be spotted in parallel on the same plate.

2. Precoated HPTLC Plates

- Glass, aluminum, or plastic-backed plates with a uniform layer of adsorbent (e.g., silica gel 60 F254).
- Particle size: 5–10 µm → ensures high resolution.
- **Plate size**: typically 10 × 10 cm or 20 × 10 cm.

3. Development Chamber (Automated or Manual)

- Saturated with mobile phase vapor before development.
- Ensures uniform solvent migration.
- Can use isocratic or gradient solvent systems.

4. Densitometer/Scanner

- Scans developed chromatograms at specific wavelengths (UV/Visible).
- Measures peak area or height → directly proportional to concentration.
- Enables quantification of separated drugs.

5. Data Station / Software

- Controls all modules: sample application, plate development, scanning.
- Performs integration, calibration, and quantitative analysis.
- Stores data for regulatory compliance and audit trails.

6. Derivatization Apparatus (Optional)

- Used to visualize colorless compounds via post-development chemical reactions.
- Example: Ninhydrin for amino acids, Dragendorff's reagent for alkaloids.

Applications of HPTLC in Pharmaceutical Analysis

A. Qualitative Analysis

- Identification of drugs by comparing Rf and spectra with standards.
- **Example:** Detecting aspirin in combined pain-relief tablets.

B. Quantitative Analysis

- Measures concentration of drug substances or impurities using peak area/height.
- Suitable for bulk drug analysis, formulations, and dissolution testing.

C. Impurity Profiling

- Detects related substances, residual solvents, and degradation products.
- Stability-indicating method for routine testing.
- **Example:** Monitoring photodegradation of nifedipine under UV exposure.

D. Drug-Excipient Compatibility

- Can determine if excipients affect the stability or recovery of drugs.
- **Example:** Separation of ibuprofen from lactose, magnesium stearate, and talc.

E. Herbal Drug Standardization

- Simultaneous identification and quantification of multiple phytoconstituents.
- Fingerprinting helps in authenticity, purity, and quality control.
- **Example:** Quantification of curcumin in turmeric extract.

F. Dissolution and Content Uniformity Testing

- Used in tablets and capsules to confirm uniform distribution of the drug.
- Validates consistency in manufacturing.

G. Bioanalytical Applications

- Though not as sensitive as LC-MS or HPLC, HPTLC can analyze biological fluids for drugs and metabolites.
- **Example:** Screening urine samples for paracetamol or caffeine.

H. Cost-effective Screening Tool

- Ideal for screening counterfeit or substandard drugs, especially in low-resource settings.
- Can test multiple samples simultaneously, making it efficient for QC labs.

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.