Noncompartmental Pharmacokinetics

Noncompartmental Pharmacokinetics (NCA): A Comprehensive Introduction

Overview

- Noncompartmental pharmacokinetics (NCA) is a method of analyzing pharmacokinetic data that does not rely on assuming specific compartmental models (like one-compartment or two-compartment models).
- Instead of defining the body as a system of connected compartments with first-order transfer rates, NCA uses mathematical and statistical tools to describe drug disposition based on the time course of drug concentrations.
- The method is model-independent, meaning it doesn't make prior assumptions about the number of compartments in the body.

Core Principles of NCA

- **Area Under the Curve (AUC):** The AUC represents the total drug exposure over time. It's calculated using the trapezoidal rule.
- **Clearance (CL):** Clearance is the volume of plasma from which the drug is completely removed per unit time. For IV administration:

$$CL = \frac{Dose_{IV}}{AUC}$$

- **Volume of Distribution at Steady State (Vss):**

$$V_{ss} = CL \times MRT$$

- **Mean Residence Time (MRT):** The average time a drug molecule spends in the body. It's derived from the area under the first moment curve (AUMC).

Statistical Moment Theory (SMT)

Introduction to SMT

- Statistical Moment Theory is a mathematical framework originally developed in the field of probability and statistics.
- In pharmacokinetics, it has been adapted to describe the distribution and elimination of drugs by analyzing concentration vs. time data using statistical moments (e.g., mean, variance).
- A moment in statistics is a quantitative measure related to the shape of a function's distribution.
- The first moment refers to the mean, and the second central moment refers to the variance. In pharmacokinetics, moments are used to describe the drug concentration-time profile.

Statistical Moment Theory is particularly useful for describing:

- **Drug exposure** (how much drug is in the body over time)
- **Drug residence time** (how long the drug stays in the body)

- It is most applicable to **linear pharmacokinetics**, where dose and concentration relationships are proportional.

Key Definitions in SMT

1. Zero-order Moment (AUC)

- The **Area Under the Curve (AUC)** is the total exposure of the body to the drug.

$$\text{AUC}_{0\to\infty} = \int_0^{\infty} C(t)\,dt$$

- Where:
  - $C(t)$is the drug concentration in plasma at time t
  - It reflects the **total amount of drug exposure** over time.

2. First-order Moment (AUMC)

- The **Area Under the Moment Curve (AUMC)** is the integral of time times the drug concentration:

$$\text{AUMC}_{0\to\infty} = \int_0^{\infty} t \cdot C(t)\,dt$$

- It reflects the **total time-weighted drug exposure** — essentially how long the drug "resides" in the body, weighted by concentration.

3. Mean Residence Time (MRT)

- MRT is the **average time** a molecule stays in the body after administration.

$$\text{MRT} = \frac{\text{AUMC}}{\text{AUC}}$$

- MRT gives a more comprehensive understanding of drug behavior than half-life alone, especially in multi-compartment or non-linear kinetics.

Why Use SMT in Pharmacokinetics?

- Statistical Moment Theory allows us to compute important pharmacokinetic parameters without assuming any specific compartmental structure.
- It's particularly useful when data is not sufficient to fit a full compartmental model or when one wants a more general description of drug kinetics.

The main pharmacokinetic parameters derived from SMT include:

- **MRT (Mean Residence Time)**
- **Clearance (CL)**

- **Volume of Distribution (Vss)**

These are calculated as follows (for IV administration):

- Clearance:

$$CL = \frac{\text{Dose}}{\text{AUC}}$$

- Volume of Distribution at Steady-State (Vss):

$$V_{ss} = CL \cdot MRT = \frac{\text{Dose} \cdot \text{AUMC}}{\text{AUC}^2}$$

- These parameters are central to understanding drug disposition using SMT.

Assumptions and Limitations

- The drug follows **linear pharmacokinetics**
- The **sampling duration** must be long enough to capture most of the drug elimination
- **Extrapolation to infinity** is often needed (especially for AUC and AUMC)

A major limitation is that SMT provides average or global properties (like MRT) rather than insights into specific phases (distribution, elimination) unless further data analysis is conducted.

MRT for Various Compartment Models

Understanding Mean Residence Time (MRT)

- MRT, as introduced earlier, is the average time a molecule remains in the body following administration.
- It's derived from AUC and AUMC.
- MRT is a global, time-based measurement that summarizes the overall retention of the drug in the system.
- The method of administration (IV vs. oral), and the number of compartments in the model affect how MRT is calculated and interpreted.

MRT for IV Bolus Administration (Single Compartment)

- For a single-compartment model with IV bolus injection:

$$C(t) = C_0 \cdot e^{-k \cdot t}$$

- Where:
  - C_0= initial concentration
  - k= elimination rate constant
- Then:

- $\text{AUC} = \frac{C_0}{k}$
- $\text{AUMC} = \frac{C_0}{k^2}$

- So,

$$\text{MRT}_{iv} = \frac{\text{AUMC}}{\text{AUC}} = \frac{1}{k}$$

- This matches the **mean lifetime of a drug molecule** in the system for a one-compartment model.

MRT for Two-Compartment Model (IV Bolus)

- In a two-compartment model, a drug distributes between central and peripheral compartments.
- The drug concentration over time is usually:

$$C(t) = Ae^{-\alpha t} + Be^{-\beta t}$$

- Where:
  - α= distribution rate constant
  - β= elimination rate constant
  - A, B= coefficients
- AUC and AUMC become more complex integrals, but can be calculated numerically or analytically. Importantly, the MRT is:

$$\text{MRT}_{iv} = \text{MRT}_{central} + \text{MRT}_{peripheral}$$

- In general:

$$\text{MRT}_{iv} = \frac{AUMC}{AUC}$$

- But the result is **greater than** $\frac{1}{\beta}$due to the time spent in peripheral compartments.
- The peripheral distribution causes **a delay in elimination**, leading to a longer MRT.

MRT for Oral Administration (Extravascular)

- With oral or other extravascular administration, absorption must be considered.
- MRT includes the time the drug takes to be absorbed **plus** the time it spends in the body.
- So, for oral administration:

$$\text{MRT}_{po} = \text{MRT}_{iv} + \text{MAT}$$

- Where:
  - **MAT** = Mean Absorption Time (Average time the drug takes to enter systemic circulation after administration)

First Hope

FREE PREVIEW

Click here to visit our website

PHARMACY PREMIUM NOTES

Complete subject and semesters are available on FirstHope app only.

Scan or click here

Click here to get the app >

Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram >

Scan or click here

@FirstHopeNotes

Instagram

@firsthope.official

Click here to follow >

YouTube channel

FirstHope-Pharmacy / Live streams

Click here to watch >

WhatsApp channel

Follow FirstHope on WhatsApp

Click here to join >

FirstHope website

firsthope.co.in

Click here to visit >

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.